


2011

Production Guide for Organic Snap Beans for Processing



NYS IPM Publication No. 132 v2



Cornell University
Cooperative Extension



New York State
Department of
Agriculture & Markets

2011 PRODUCTION GUIDE FOR ORGANIC BEANS FOR PROCESSING

Coordinating Editor

Abby Seaman* (NYS IPM Program)

Contributors and Resources

George Abawi (Cornell University, Department of Plant Pathology)

Thomas Bjorkman (Cornell University, Vegetable Crop Physiology)

Ann Cobb (Cornell University, Department of Plant Pathology)

Helene Dillard (Cornell University, Department of Plant Pathology)

Vern Grubinger (University of Vermont, Vegetable and Berry Specialist)

Beth Gugino (The Pennsylvania State University, Department of Plant Pathology)

Robert Hadad (Cornell Cooperative Extension Vegetable Specialist)

Julie Kikkert (Cornell Cooperative Extension, Department of Horticulture)

Michael Helms* (Pesticide Management Education Program)

Margaret T. McGrath (Long Island Horticultural Research and Extension Center, Department of Plant Pathology)

Charles L. Mohler (Cornell University, Weed Ecology)

Brian Nault* (Cornell University, Department of Entomology)

Anu Rangarajan (Cornell Small Farms Program, Department of Horticulture)

Thomas A. Zitter (Cornell University, Plant Pathology)

**Pesticide Information and Regulatory Compliance*

Staff Writers

Mary Kirkwyland and Elizabeth Thomas (New York State IPM Program)

Special Appreciation

Format based on the Integrated Crop and Pest Management Guidelines for Commercial Vegetable Production. Content Editors Stephen Reiners and Curtis H. Petzoldt, with numerous Discipline Editors.

Funded in part by the New York State Department of Agriculture and Markets

The information in this guide reflects the current authors' best effort to interpret a complex body of scientific research, and to translate this into practical management options. Following the guidance provided in this guide does not assure compliance with any applicable law, rule, regulation or standard, or the achievement of particular discharge levels from agricultural land.

Every effort has been made to provide correct, complete, and up-to-date pest management information for New York State at the time this publication was released for printing (February 2011). Changes in pesticide registrations and regulations, occurring after publication are available in county Cornell Cooperative Extension offices or from the Pesticide Management Education Program web site (<http://pmep.cce.cornell.edu>). Trade names used herein are for convenience only. No endorsement of products is intended, nor is criticism of unnamed products implied.

This guide is not a substitute for pesticide labeling. Always read the product label before applying any pesticide.

Updates and additions to this guide are available at http://www.nysipm.cornell.edu/organic_guide. Please submit comments or suggested changes for these guides to organicguides@gmail.com.

TABLE OF CONTENTS

1. General Organic Management Practices	3
2. Soil Health.....	3
3. Cover Crops.....	2
4. Field Selection.....	4
5. Weed Management	6
6. Recommended Varieties	6
7. Planting Methods.....	7
8. Crop & Soil Nutrient Management	7
9. Harvesting.....	11
10. Using Organic Pesticides	11
11. Disease Management.....	13
11.1 Sclerotinia White Mold, <i>Sclerotinia sclerotiorum</i>	15
11.2 Botrytis Gray Mold, <i>Botrytis cinerea</i>	16
11.3 Pod-flecking complex (PFC), <i>Alternaria alternata</i> and <i>Plectosporium tabacinum</i>	18
11.4 Root Rot and Damping-Off.....	19
11.5 Bacterial Diseases.....	21
11.6 Virus Diseases.....	23
11.7 Bean Rust, <i>Uromyces appendiculatus</i>	24
12. Root-Lesion Nematode, <i>Pratylenchus penetrans</i>.....	25
13. Insect Management	26
13.1 Mexican Bean Beetle.....	28
13.2 Potato Leafhopper (PLH), <i>Empoasca fabae</i>	30
13.3 Seedcorn Maggot, <i>Delia platura</i>	31
13.4 European Corn Borer (ECB), <i>Ostrinia nubilalis</i>	32
13.5 Two-Spotted Spider Mite, <i>Tetranychus urticae</i>	33
13.6 Tarnished Plant Bug (TPB), <i>Lygus lineolaris</i>	34
14. Slugs.....	36
15. Pesticides & Abbreviations Mentioned in this Publication	37
16. References	38
17. World Wide Web Links.....	39

INTRODUCTION

This guide for organic production of snap beans provides an outline of cultural and pest management practices and includes topics that have an impact on improving plant health and reducing pest problems. It is divided into sections, but the interrelated quality of organic cropping systems makes each section relevant to the others.

The guide attempts to compile the most current information available, but acknowledges that effective means of control are not available for some pests. More research on growing crops organically is needed, especially in the area of pest management. Future revisions will incorporate new information, providing organic growers with a complete set of useful practices to help them achieve success.

This guide uses the term Integrated Pest Management (IPM), which like organic production, emphasizes cultural, biological, and mechanical practices to minimize pest outbreaks. With limited pest control products available for use in many organic production systems, an integrated approach to pest management is essential. IPM techniques such as identifying and assessing pest populations, keeping accurate pest history records, selecting the proper site, and preventing pest outbreaks through use of crop rotation, resistant varieties and biological controls are important to producing a high quality crop.

1. GENERAL ORGANIC MANAGEMENT PRACTICES

1.1 Organic Certification

To use a certified organic label, farming operations that gross more than \$5,000 per year in organic products must be certified by a U.S. Department of Agriculture National Organic Program (NOP) accredited certifying agency. The choice of certifier may be dictated by the processor or by the target market. [A list of accredited certifiers](#) (Link 6) operating in New York can be found on the New York State Department of Agriculture and Markets [Organic Farming Resource Center web page](#) (Link 7). See more certification and regulatory details under Section 4.1: *Certification Requirements* and Section 10: *Using Organic Pesticides*.

1.2 Organic Farm Plan

An organic farm plan is central to the certification process. The farm plan describes production, handling, and record-keeping systems, and demonstrates to certifiers an understanding of organic practices for a specific crop. The process of developing the plan can be very valuable in terms of anticipating potential issues and challenges, and fosters thinking of the farm as a whole system. Soil,

nutrient, pest, and weed management are all interrelated on organic farms and must be managed in concert for success. Certifying organizations may be able to provide a template for the farm plan. The following description of the farm plan is from the NOP web site:

The Organic Food Production Act of 1990 (OFPA or Act) requires that all crop, wild crop, livestock, and handling operations requiring certification submit an organic system plan to their certifying agent and, where applicable, the State Organic Program (SOP). The organic system plan is a detailed description of how an operation will achieve, document, and sustain compliance with all applicable provisions in the OFPA and these regulations. The certifying agent must concur that the proposed organic system plan fulfills the requirements of subpart C, and any subsequent modification of the organic plan by the producer or handler must receive the approval of the certifying agent.

More details may be found at the Agricultural Marketing Service's [National Organic Program website](#) (Link 8). The [National Sustainable Agriculture Information Service](#), (formerly ATTRA), has produced a guide to organic certification that includes templates for developing an organic farm plan (Link 10). The [Rodale Institute](#) has also developed resources for transitioning to organic and developing an organic farm plan (Link 11).

2. SOIL HEALTH

Healthy soil is the basis of organic farming. Regular additions of organic matter in the form of cover crops, compost, or manure create a soil that is biologically active, with good structure and capacity to hold nutrients and water (note that any raw manure applications should occur at least 120 days before harvest). Decomposing plant materials will activate a diverse pool of microbes, including those that break down organic matter into plant-available nutrients as well as others that compete with plant pathogens on the root surface.

Rotating between crop families can help prevent the buildup of diseases that overwinter in the soil. Rotation with a grain crop, preferably a sod that will be in place for one or more seasons, deprives disease-causing organisms of a host, and also contributes to a healthy soil structure that promotes vigorous plant growth. The same practices are effective for preventing the buildup of root damaging nematodes in the soil, but keep in mind that certain grain crops are also hosts for some nematode species. Rotating between crops with late and early season planting dates can help prevent the buildup of weed populations. Organic growers must attend to the connection between soil, nutrients, pests, and weeds to succeed. An excellent resource for additional information on soils and soil health is [Building Soils for Better Crops](#) by Fred Magdoff and Harold Van Es, 2000 (Link 13). For additional

information, refer to the [Cornell Soil Health website](#) (Link 14).

3. COVER CROPS

Unlike cash crops, which are grown for immediate economic benefit, cover crops are grown for their valuable effect on soil properties and on subsequent cash crops. Cover crops help maintain soil organic matter, improve soil tilth, prevent erosion and assist in nutrient management. They can also contribute to weed management, increase water infiltration, maintain populations of beneficial fungi, and may help control insects, diseases and nematodes. To be effective, cover crops should be treated as any other valuable crop on the farm, with their cultural requirements carefully considered including their cultural requirements, life span, mowing recommendations, incorporation methods, and susceptibility, tolerance, or antagonism to root pathogens and other pests. Some cover crops and cash crops share susceptibility to certain pathogens and nematodes. Careful planning and monitoring is required when choosing a cover crop sequence to avoid increasing pest problems in subsequent cash crops. See Tables 3.1 for more information on specific cover crops and Section 8: *Crop and Soil Nutrient Management* for more information about how cover crops fit into a nutrient management plan.

A certified organic farmer is required to plant certified organic cover crop seed. If, after contacting at least three suppliers, organic seed is not available, then the certifier may allow conventional seed to be used. Suppliers should provide a purity test for cover crop seed. Always inspect the seed for contamination with weed seeds and return if it is not clean. Cover crop seed is a common route for introduction of new weed species onto farms.

3.1 Goals and Timing for Cover Crops

Adding cover crops regularly to the crop rotation plan can result in increased yields of the subsequent cash crop. Goals should be established for choosing a cover crop; for example, the crop can add nitrogen, smother weeds, or break a pest cycle. The cover crop might best achieve some of these goals if it is in place for the entire growing season. If this is impractical, a compromise might be to grow the cover crop between summer cash crops. Allow two or more weeks between cover crop incorporation and cash crop seeding to permit decomposition of the cover crop, which will improve the seedbed and help avoid any unwanted allelopathic effects on the next cash crop. Another option is to overlap the cover crop and the cash crop life cycles by overseeding, interseeding or intercropping the cover crop between cash crop rows at final cultivation. An excellent resource for

determining the best cover crop for your situation is [Northeast Cover Crop Handbook](#), by Marianne Sarrantonio (Reference 11) or the [Cornell online decision tool](#) to match goals, season, and cover crop (Link 12).

Leaving cover crop residue to remain on the soil surface might make it easier to fit into a crop rotation and will help to conserve soil moisture, but some of the nitrogen contained in the residue will be lost to the atmosphere, and total organic matter added to the soil will be reduced. Turning under the cover crop will speed up the decomposition and nitrogen release from the crop residue.

3.2 Legumes Cover Crops

Legume cover crops should be avoided before beans because many are closely related to beans and share pests.

3.3 Non-Legume Cover Crops

Barley, rye grain, rye grass, Sudangrass, wheat, oats, and other grain crops left on the surface or plowed under as green manures or dry residue in the spring are beneficial because these plants take up nitrogen that otherwise might be leached from the soil, and release it back to the soil as they decompose. If incorporated, allow two weeks or more for decomposition prior to planting to avoid the negative impact on stand establishment from actively decomposing material. Three weeks might not be enough if soils are very cold. In wet years, the presence of cover crop residues may increase slug damage and infections by fungal pathogens such as *Pythium* and *Rhizoctonia*, affecting stand establishment

3.4 Biofumigant Cover Crops

Certain cover crops have been shown to inhibit weeds, pathogens, and nematodes by releasing toxic volatile chemicals when tilled into the soil as green manures and degraded by microbes or when cells are broken down by finely chopping. Degradation is quickest when soil is warm and moist. These biofumigant cover crops include Sudangrass, sorghum-sudangrasses, and many in the brassica family. Varieties of mustard and arugula developed with high glucosinolate levels that maximize biofumigant activity have been commercialized (e.g. Caliente brands 199 and Nemat).

Attend to the cultural requirements of the cover crops to maximize growth. Fertilizer applied to the cover crops will be taken up and then returned to the soil for use by the cash crop after the cover crop is incorporated. Biofumigant cover crops like mustard should be allowed to grow to their full size, normally several weeks after flowering starts, but incorporated before the seeds become brown and hard indicating they are mature. To minimize loss of biofumigant, finely chop the tissue early in the day when temperatures are

ORGANIC BEAN PRODUCTION

low. Incorporate immediately by tilling, preferably with a second tractor following the chopper. Lightly seal the soil surface using a culti-packer and/or 1/2 inch of irrigation or rain water to help trap the volatiles and prolong their persistence in the soil. Wait at least two weeks before planting a subsequent crop to reduce the potential for the breakdown products to harm the crop, also known as phytotoxicity. Scratching the soil surface before planting will release remaining biofumigant. This biofumigant effect is not predictable or consistent. The levels of the active compounds

and suppressiveness can vary by season, cover crop variety, maturity at incorporation, amount of biomass, fineness of chopping, how quickly the tissue is incorporated, soil microbial diversity, soil tilth, and microbe population density.

[Cover Crops for Vegetable Growers: Decision Tool](#) (Link 12).
[Northeast Cover Crops Handbook](#) (Reference 11).
[Cover Crops for Vegetable Production in the Northeast](#) (Ref. 17).
[Crop Rotation on Organic Farms: A Planning Manual](#) (Link 14a).

Table 3.1. Non-Leguminous Cover Crops: Cultural Requirements and Crop Benefits

SPECIES	PLANTING DATES	LIFE CYCLE	COLD HARDINESS ZONE	HEAT	DROUGHT	SHADE	PH PREFERENCE	SOIL TYPE PREFERENCE	SEEDING (LB/A)	COMMENTS
				--TOLERANCES--						
Brassicas e.g. mustards, rapeseed	April or late August-early Sept.	Annual / Biennial **	6-8	4	6	NI	5.3-6.8	Loam to clay	5-12	+Good dual purpose cover & forage +Establishes quickly in cool weather +Biofumigant properties
Buckwheat	Late spring-summer	Summer annual **	NFT	7-8	4	6	5.0-7.0	Most	35-134	+Rapid grower (warm season) +Good catch or smother crop +Good short-term soil improver for poor soils
Cereal Rye	August-early October	Winter annual	3	6	8	7	5.0-7.0	Sandy to clay loams	60-200	+Most cold-tolerant cover crop +Excellent allelopathic weed control +Good catch crop +Rapid germination & growth +Temporary N tie-up when turned under
Fine Fescues	Mid March-mid-May OR late Aug.-late Sept.	Long-lived perennial	4	3-5	7-9	7-8	5.3-7.5 (red) 5.0-6.0 (hard)	Most	16-100	+Very good low-maintenance permanent cover, especially in infertile, acid, droughty &/or shady sites
Oats	Mid-Sept-early October	Summer annual**	8	4	4	4	5.0-6.5	Silt & clay loams	110	+Rapid growth +Ideal quick cover and nurse crop
Ryegrasses	August-early Sept.	Winter annual (AR)/ Short-lived perennial (PR)	6 (AR) 4 (PR)	4	3	7 (AR) 5 (PR)	6.0-7.0	Most	14-35	+Temporary N tie-up when turned under +Rapid growth +Good catch crop +Heavy N & moisture users
Sorghum-Sudangrass	Late spring-summer	Summer Annual **	NFT	9	8	NI	Near neutral	NI	10-36	+Tremendous biomass producers in hot weather +Good catch or smother crop +Biofumigant properties

NI-No Information, NFT-No Frost Tolerance. Drought, Heat, Shade Tolerance Ratings: 1-2=low, 3-5=moderate, 6-8=high, 9-10=very high. . **Winter killed.

AR=Annual Rye, PR=Perennial Rye.

Reprinted with permission from Rodale Institute®, www.rodaleinstitute.org, M. Sarrantonio. 1994. Northeast Cover Crop Handbook. (Reference 11).

4. FIELD SELECTION

For organic production, give priority to fields with excellent soil tilth, high organic matter, good drainage and airflow. Beans do not thrive in wet soil.

4.1 Certifying Requirements

Certifying agencies have requirements that affect field selection. Fields cannot be treated with prohibited products for three years prior to the harvest of a certified organic crop. Adequate buffer zones are required between certified organic and conventionally grown crops. Buffer zones must be a barrier, such as a diversion ditch or dense hedgerow, or be a distance large enough to prevent drift of prohibited materials onto certified organic fields. Determining what buffer zone is needed will vary depending on equipment used on adjacent non-certified land. For example, use of high-pressure spray equipment or aerial pesticide applications in adjacent fields will increase the buffer zone size. Pollen from genetically engineered crops can also be a contaminant. An organic crop should not be grown near a genetically engineered crop of the same species. Check with your certifier for specific buffer requirements. These buffers commonly range between 20 to 250 feet depending on adjacent field practices.

4.2 Crop Rotation Plan

A careful crop rotation plan is the cornerstone of organic crop production because it allows the grower to improve soil quality and proactively manage pests. Although growing a wide range of crops complicates the crop rotation planning process, it ensures diversity in crop residues in the soil, and a greater variety of beneficial soil organisms. Individual organic farms vary widely in the crops grown and their ultimate goals, but some general rules apply to all organic farms regarding crop rotation. Rotating individual fields away from crops within the same family is critical and can help minimize crop-specific disease and non-mobile insect pests that persist in the soil or overwinter in the field or field borders. Pests that are persistent in the soil, have a wide host range, or are wind-borne, will be difficult to control through crop rotation. Conversely, the more host specific, non-mobile, and short-lived a pest is, the greater the ability to control it through crop rotation. The amount of time required for a crop rotation is based on the particular pest and its severity. Some particularly difficult pests may require a period of fallow. See specific recommendations in the disease and insect sections of this guide (Sections 11, 12, 13). Partitioning the farm into management units will help to organize crop rotations and ensure that all parts of the farm have sufficient breaks from each type of crop.

A well-planned crop rotation is key to weed management. Short season crops such as lettuce and spinach are harvested before many weeds go to seed, whereas vining cucurbits, with their limited cultivation time and long growing season, allow weeds to go to seed before harvest. Including short season crops in the rotation will help to reduce weed populations provided the field is cleaned up promptly after harvest. Other weed reducing rotation strategies include growing mulched crops, competitive cash crops, short-lived cover crops, or crops that can be intensively cultivated. Individual weed species emerge and mature at different times of the year, therefore alternating between spring, summer, and fall planted crops helps to interrupt weed life cycles.

Cash and cover crop sequences should also take into account the nutrient needs of different crops and the response of weeds to high nutrient levels. High soil phosphorus and potassium levels can exacerbate problem weed species. A cropping sequence that alternates crops with high and low nutrient requirements can help keep nutrients in balance. The crop with low nutrient requirements can help use up nutrients from a previous heavy feeder. A fall planting of a non-legume cover crop will help hold nitrogen not used by the previous crop. This nitrogen is then released when the cover crop is incorporated in the spring. See Section 5: *Weed Management*, and Section 3: *Cover Crops* for more specifics.

Rotating crops that produce abundant organic matter, such as hay crop and grain-legume cover crops, with ones that produce less, such as vegetables, will help to sustain organic matter levels and promote good soil tilth (see Section 2: *Soil Health* and Section 8: *Crop and Soil Nutrient Management*). Beans generally have a lower nutrient requirement (Table 4.2.1). Growing a cover crop, preferably one that includes a legume (unless the field has a history of *Pythium* or *Rhizoctonia* problems), prior to or after a bean crop, will help to renew soil nutrients, improve soil structure, and diversify soil organisms. Deep-rooted crops in the rotation to help break up compacted soil layers.

Table 4.2.1 Crops Nutrient Requirements

Crop	Nutrient Needs		
	Lower	Medium	Higher
bean		cucumber	broccoli
beet		eggplant	cabbage
carrot		brassica greens	cauliflower
herbs		pepper	corn
pea		pumpkin	lettuce
radish		spinach	potato
		chard	tomato
		squash	
		winter squash	

From NRAES publication *Crop Rotation on Organic Farms: A Planning Manual*. Charles L. Mohler and Sue Ellen Johnson, editors, (Reference 14a).

ORGANIC BEAN PRODUCTION

Crop information specific to Beans

Beans should only be grown in fields that have had no legumes for 3 years. Legumes including soybean, clovers, alfalfa and hairy vetch are hosts for many soil-borne fungal pathogens and should be avoided in fields with severe root

rot problems. A good rotation helps reduce the incidence of foliar diseases and lowers the population of plant pathogens that cause root rot. Corn and cereal grains are excellent rotation crops to reduce root rot problems because they are not hosts for root rot pathogens of vegetable crops.

Table 4.2.2 Potential Interactions of Crops Grown in Rotation with Beans

Crops in Rotation	Potential Rotation Effects	Comments
Lettuce, potato, tomato, other legumes, crucifers, or cucurbits	Increase Sclerotinia	Avoid growing these crops prior to or after snap beans to reduce the buildup of Sclerotinia in the soil. Grow grains or corn for several years to help reduce <i>Sclerotinia</i> .
Bean, tomato, cucumber, buckwheat, aromatic herbs	Decrease Clubroot	Clubroot in brassicas has declined more quickly in fields where tomato, cucumber, snap bean and buckwheat have been grown. Aromatic perennial herbs such as summer savory, peppermint, or garden thyme helps to reduce clubroot when grown for 2 to 3 consecutive years.
Soybean, dry bean	Increases Bacterial blight (<i>Xanthomonas campestris</i>)	<i>Xanthomonas campestris</i> is found in legumes as well as some crucifers and weeds in the mustard family.
Soybean, dry bean	Increases Soybean cyst nematode	Soybean cyst nematode <i>heterodera glycines</i> increases to high densities on snap beans through the snap bean crop is rarely affected.
Snap bean, soybean, dry bean, alfalfa	Increase Fusarium, Pythium, and Sclerotinia	Do not plant these legumes prior to or for several years after snap beans to prevent a buildup of these soilborne diseases.

Excerpt from Appendix 2 of Crop Rotation on Organic Farms: A Planning Manual. Charles L. Mohler and Sue Ellen Johnson, editors. (Reference 14a)

4.3 Pest History

Knowledge about the pest history for each field to plan a successful cropping strategy. Germination may be reduced in fields with a history of *Pythium* or *Rhizoctonia*. Avoid fields that contain heavy infestations of perennial weeds such as nutsedge, bindweed, and quackgrass as these weeds are particularly difficult to control. One or more years focusing on weed population reduction using cultivated fallow and cover cropping may be needed before organic crops can be successfully grown in those fields. Susceptible crops should not be grown in fields with a history of *Sclerotinia* white mold without a rotation of several years to sweet corn or grain crops. If there is a field history of white mold, beans should not be preceded by tomato, potato, lettuce, crucifer crops, or bean (including soybean). Treat with Contans™ to reduce fungal sclerotia in the soil immediately after an infected crop is harvested.

If possible, beans should not be grown in fields with a history of root rot problems, but if there is no choice, plant as late as possible when the soil has warmed.

Root maggots prefer to lay eggs in soil with fresh organic matter. Incorporate cover crop residues 2-3 weeks before planting to allow time for decomposition.

Snap bean is a host for root-lesion nematode, *Pratylenchus penetrans*, and therefore it is important to know whether or not lesion nematode is present in the field in order to develop long-term crop rotations and cropping sequences that either reduce the populations in heavily infested fields or minimize their increase in fields that have no to low infestation levels. Refer to Section 12 for more information on nematodes.

4.4 Soil and Air Drainage

Most fungal and bacterial pathogens need free water on the plant tissue or high humidity for several hours in order to infect. Any practice that promotes leaf drying or drainage of excess water from the root zone will minimize favorable conditions for infection and disease development. Fields with poor air movement, such as those surrounded by hedgerows or woods, result in leaves staying wet. Plant rows parallel to the prevailing winds, which is typically in an east-west direction and avoid overcrowding by using wide row spacing to promote drying of the soil and reduce moisture in the plant canopy.

5. WEED MANAGEMENT

Weed management can be one of the biggest challenges on organic farms, especially during the transition and the first several years of organic production. To be successful, weed management on organic farms must take an integrated approach that includes crop rotation, cover cropping, cultivation, and planting design, based on an understanding of dominant weed biology and ecology of dominant weed species. A multi-year approach that includes strategies for controlling problem weed species in a sequence of crops will generally be more successful than attempting to manage each year's weeds as they appear. Relying on cultivation alone to manage weeds in an organic system is a recipe for disaster.

Management plans should focus on the most challenging and potentially yield-limiting weed species in each field. Be sure, however, to emphasize options that do not increase other species that are present. Alternating between early and late-planted crops, and short and long season crops in the rotation can help minimize buildup of a particular weed or group of weeds with similar life cycles or growth habits, and will also provide windows for a variety of cover crops.

5.1 Record Keeping

Scout and develop a written inventory of weed species and their severity for each field. Accurate identification of weeds is essential. Weed fact sheets provide a good color reference for common weed identification. See Cornell [weed ecology](#) and Rutgers [weed gallery](#) websites (Link 16 and 17).

5.2 Weed Management Methods

Planting and cultivation equipment should be set up on the same number of rows to minimize crop losses and damage to crop roots during cultivation. It may be necessary to purchase specialized equipment to successfully control weeds in some crops.

Begin blind cultivation with a tine weeder, or flexible harrow, just before ground crack, when weeds are at white thread stage. Beans are very susceptible to breakage when they are in the "crook" stage – from just before ground crack until the seed leaves are unfolded and horizontal. Avoid tine weeding during this period. Correct cultivation depth is 2/3rd of seeding depth. Note, however, that penetration will vary with soil conditions and you must avoid hitting the seed with the weeder in soft spots. Effective tine weeding is an art that requires adjustment of the weeder to obtain good weed control without harming the crop. Examples of tine weeders are the Einbock, Lely and the Kovar. The tines on various brands and models of harrows differ in flexibility. Tines that are too stiff can break bean stems. Tines with a 70 to 80

degree bend work well for beans as they hook out grassy weeds without pulling out the beans, which have a taproot. Tines with a 45-degree bend can also be used effectively.

After bean emergence, make two more passes using a tine weeder at about 5 to 7 day intervals depending on weed growth. Tine weeders work best on very small weeds. The final tine weeding can be more aggressive (faster and deeper) than the pre-emergence weeding or early post-emergence weeding. Test settings on a small area and adjust.

The flex tine harrow is effective with weeds in the row, but when crop reaches the early first trifoliolate stage, the tine weeders may cause damage to the crop. At this point, a row crop cultivator can be used to control weeds between the rows. Adjust row crop cultivator for close and shallow cultivation. Cultivate to minimize disturbing the soil and uprooting rocks that will cause problems with the harvester. Perennial weeds will require deeper cultivation. Bean varieties that grow higher off the ground will minimize rocks in the harvester. Using a rolling cultivator is another option.

For a field with persistent perennial weeds such as field bindweed, several tactics might be needed to reduce competition: tillage to break up rhizomes, one or more short fallow periods to exhaust rhizome reserves, and planting a crop that requires multiple cultivations.

Resources

[Steel in the Field](#) (Link 15).

[Cornell Weed Ecology website](#) (Link 16).

[Rutgers University, New Jersey Weed Gallery](#) (Link 17).

[Univ. of Vermont videos on cultivation and cover cropping](#) (Link 18).

[ATTRA Principles of Sustainable Weed Mgt. for Croplands](#) (Link 19).

[Cultivation Tools for Mechanical Weed Control in Vegetables](#) (Link 20)

6. RECOMMENDED VARIETIES

Variety selection is important both for the horticultural characteristics specified by the processor and the pest resistance profile that will be the foundation of a pest management program. Collaborate with processors on varieties, choosing those with some level of disease resistance if possible. Cornell research on developing CMV-resistant snap bean varieties is ongoing.

A certified organic farmer is required to plant certified organic seed. If, after contacting at least three suppliers, organic seed is not available for a particular variety, then the certifier may allow untreated conventional seed to be used.

6.1 Varieties Currently Grown for Processing in NY

Large Sieve Green	Wax Bean Type
Hystyle	Goldmine
Igloo	Serin
Spartacus	703
Summit	Whole Bean Type
Titan	Banga
Venture	Masai
Weapon	Pix
5402	PLS75
Romano Type	1267
Tapia	

6.2 Varieties Identified as Candidates for Organic Production*

Sieve Green	Whole Bean Green
Caprice	Banga
Diplomat	Celtic
Ducato	
Tesco	

*Suggested for trial use only until more information is available

7. PLANTING METHODS

Recommended earliest planting date for untreated snap bean seed is June 1. The crop matures in 50 to 60 days, depending on the specific variety and desired pod size. Only western-grown, certified seed should be planted. Optimal germination of snap bean seed occurs at soil temperatures of 75° to 80°F. The minimum temperature at which snap bean germination will occur is 55° to 60°F. Plant rows in an east-west direction if possible and use wide row spacing, 36 inches, 5-7 plants/foot, to promote drying of the soil, increase air circulation, reduce moisture in the plant canopy and reduce risk of foliar diseases. Bean seed is sensitive to chilling during the initial stage of germination. If the soil is cold at this time, permanent damage may occur.

A good rotation helps reduce the incidence of foliar diseases and lowers the population of plant pathogens that cause root rot. Corn and cereal grains are excellent rotation crops. Planting on raised beds or ridges will help reduce root rot severity because the soil will be warmer and drier than the unridged soil. To allow for adequate aeration and drainage of excess moisture, avoid compacting the soil.

A water deficiency resulting from a lack of soil moisture or excessive transpiration can lead to deformed or pithy snap bean pods. Both yield and quality can be increased by irrigation before bloom and during pod enlargement if there is moisture stress. Irrigation during bloom with irrigation guns that produce large droplets is not advised because blossoms can be knocked off the plant.

8. CROP & SOIL NUTRIENT MANAGEMENT

To produce a healthy crop, soluble nutrients must be available from the soil in amounts that meet the minimum requirements for the whole plant. The total nutrient needs of a crop are much higher than just the nutrients that are removed from the field when that crop is harvested. All of the roots, stems, leaves and other plant parts require nutrients at specific times during plant growth and development. The challenge in organic systems is balancing soil fertility to supply these required plant nutrients at a time and at sufficient levels to support healthy plant growth. Restrictions in any one of the needed nutrients will slow growth and can reduce crop quality and yields.

Organic growers often speak of feeding the soil rather than feeding the plant. A more accurate statement is that organic growers focus their fertility program on feeding soil microorganisms rather than the plant. Soil microbes decompose organic matter to release nutrients and convert organic matter to more stable forms such as humus. This breakdown of soil organic matter occurs throughout the growing season, depending on soil temperatures, water availability and soil quality. The released nutrients are then held on soil particles or humus making them available to crops or cover crops for plant growth. Amending soils with compost, cover crops, or crop residues also provides a food source for soil microorganisms and when turned into the soil, starts the nutrient cycle again.

During the transition years and the early years of organic production, soil amendment with composts or animal manure can be a productive strategy for building organic matter, biological activity and soil nutrient levels. This practice of heavy compost or manure use is not, however, sustainable in the long-term. If composts and manures are applied in the amounts required to meet the nitrogen needs of the crop, phosphorous may be added at higher levels than required by most vegetable crops. This excess phosphorous will gradually build up to excessive levels, increasing risks of water pollution or invigorating weeds like purslane and pigweed. A more sustainable, long-term approach is to rely more on legume cover crops to supply most of the nitrogen needed by the crop and use grain or grass cover crops to capture excess nitrogen released from organic matter at the end of the season to minimize nitrogen losses to leaching. See Section 3: *Cover Crops*. When these cover crops are incorporated into the soil, their nitrogen, as well as carbon, feeds soil microorganisms, supporting the nutrient cycle. Harvesting alfalfa hay from the field for several years can reduce high phosphorus and potassium levels.

The primary challenge in organic systems is synchronizing nutrient release from organic sources, particularly nitrogen,

with the crop requirements. In cool soils, microorganisms are less active, and nutrient release may be too slow to meet the crop needs. Once the soil warms, nutrient release may exceed crop needs. In a long-term organic nutrient management approach, most of the required crop nutrients would be in place as organic matter before the growing season starts. Nutrients required by the crop in the early season can be supplemented by highly soluble organic amendments such as poultry manure composts or organically approved bagged fertilizer products (see Tables 8.2.4 to 8.2.6). These products can be expensive so are most efficiently used if banded at planting. The National Organic Standards Board states that no more than 20% of total N can be applied as Chilean nitrate. Be sure to confirm the practice with your organic certifier prior to field application.

Regular soil testing helps monitor nutrient levels, in particular phosphorus (P) and potassium (K). Choose a reputable soil-testing lab (Table 8.0.1) and use it consistently to avoid discrepancies caused by different soil extraction methods. Maintaining a soil pH between 6.3 and 6.8 will maximize the availability of all nutrients to plants.

Table 8.0.1 Nutrient Testing Laboratories

TESTING LABORATORY	SOIL	COMPOST/ MANURE	FORAGE	LINK
Cornell Soil Health Lab	x			14
Agri Analysis, Inc.		x		22
A&L Eastern Ag Laboratories, Inc.	x	x		23
Penn State Ag Analytical Services Lab.	x	x		24
University of Massachusetts	x	x		26
The Agro One Lab			x	25

Develop a plan for estimating the amount of nutrients that will be released from soil organic matter, cover crops, compost, and manure. A strategy for doing this is outlined in Section 8.2: *Preparing an Organic Nutrient Budget*.

8.1 Fertility

Recommendations from the Cornell Integrated Crop and Pest Management Guidelines indicate that a bean crop requires 40 lbs. of available nitrogen (N), 80 lbs. of phosphorus (P), and 60 lbs. of potassium (K) per acre. These levels are based on the total needs of the whole plant and assume the use of synthetic fertilizers. Farmer and research experience suggests that lower levels may be adequate in organic systems. See Table 8.2.2 for the recommended rates of P and K based on soil test results. Nitrogen is not included because levels of available N change in response to soil temperature and moisture, N mineralization potential, and

leaching. As many of the nutrients as possible should come from cover crop, manure, and compost additions in previous seasons.

The source of these nutrients depends on soil type and historic soil management. Some soils are naturally high in P and K, or have a history of manure applications that have resulted in elevated levels. Additional plant available nutrients are supplied by decomposed soil organic matter or through specific soluble nutrient amendments applied during the growing season in organically managed systems. Many types of organic fertilizers are available to supplement the nutrients supplied by the soil. **ALWAYS check with your certifier before using any product to be sure it is approved.**

8.2 Preparing an Organic Nutrient Budget

Insuring an adequate supply of nutrients when the crop needs them requires careful planning. Developing an organic nitrogen budget can help estimate the amount of nutrients released by various organic amendments as well as native soil organic matter. Table 8.2.3 estimates common nutrient content in animal manures; however actual compost and manure nutrient content should be tested just prior to application. Analysis of other amendments, as well as cover crops, can be estimated using published values (see Tables 8.2.4 to 8.2.6 and 3.1 for examples). Keeping records of these nutrient inputs and subsequent crop performance will help evaluate if the plan is providing adequate fertility during the season to meet production goals.

Remember that with a long-term approach to organic soil fertility, the N mineralization rates of the soil will increase. This means that more N will be available from organic amendments because of increased soil microbial activity and diversity. Feeding these organisms different types of organic matter is essential to building this type of diverse biological community and ensuring long-term organic soil and crop productivity. Consider submitting soil samples for a [Cornell Soil Health Test](#) (Link 14). This test includes an estimate of nitrogen mineralization rate, which indicates the potential for release of N from soil organic matter. Testing soils over time can be useful for monitoring changes in nitrogen mineralization rate during the transition, and over time, in organic production.

Estimating total nutrient release from the soil and comparing it with soil test results and recommendations requires record-keeping and some simple calculations. Table 8.2.1 below can be used as a worksheet for calculating nutrients supplied by the soil compared to the total crop needs.

Table 8.2.1 Calculating Nutrient Credits and Needs.

	Nitrogen (N) lbs/A	Phosphate (P ₂ O ₅) lbs/A	Potash (K ₂ O)lbs/A
1. Total crop nutrient needs			
2. Recommendations based on soil test	Not provided		
3. Credits			
a. Soil organic matter		---	---
b. Manure			
c. Compost			
d. Prior cover crop			
4. Total credits:			
5. Additional needs (2-4=)			

Line 1. Total Crop Nutrient Needs: Research indicates that an average bean crop requires 40 lbs. of available nitrogen (N), 80 lbs. of phosphorus (P), and 60 lbs. of potassium (K) per acre to support a medium to high yield crop (see Section 8.1 *Fertility* above).

Line 2. Recommendations Based on Soil Test: Use Table 8.2.2 to determine the amount of P and K needed based on soil test results.

Table 8.2.2 Recommended Amounts of Phosphorus and Potassium for Beans Based on Soil Tests

	Soil Phosphorus Level			Soil Potassium Level		
	low	med	high	low	med	high
	P ₂ O ₅ lbs/A			K ₂ O lbs/A		
Total nutrient recommendation	80	60	40	60	40	20

Line 3a. Soil Organic Matter: Using the values from your soil test, estimate that 20 lbs. of nitrogen will be released from each percent organic matter in the soil. For example, a soil that has 3% organic matter could be expected to provide 60 pounds of N per acre.

Line 3b. Manure: Assume that applied manure will release N for 3 years. Based on the test of total N in any manure applied, estimate that 50% is available in the first year, and then 50% of the remaining is released in each of the next two years. For an application rate of 100 lbs. of N as manure, 50 lbs. would be available the first year, 25 lbs. in year 2, and 12.5 lbs. in year 3. The NOP rules allow manure applications 120 days or more before harvest of a vegetable crop, but your farm certifier may have a more restrictive policy.

Line 3c. Compost: Estimate that between 10 and 25% of the N contained in compost will be available the first year. Compost maturity will influence how much N is available. If the material is immature, more of the N may be available to the crop in the first year. A word of caution: Using compost to provide for a crop’s nutrient needs is not generally a financially viable strategy. The total volume needed can be very expensive for the units of N available to the crop, especially if trucking is required. Most stable composts should be considered as soil conditioners, improving soil health, microbial diversity, tilth, and nutrient retaining capacity. Any compost applied on organic farms must be approved for use by your farm certifier. Compost generated on the farm must follow an approved process outlined by your certifier.

Line 3d. Cover Crops: Legume cover crops are a source of nitrogen but are not recommended prior to a bean crop.

Line 4. Total Credits: Add together the various N values from the organic matter, compost, and cover crops to estimate the N supplying potential of the soil (see example below). There is no guarantee that these amounts will actually be available in the season, since soil temperatures, water, and crop physiology all impact the release and uptake of these soil nutrients

ORGANIC SNAP BEAN PRODUCTION

Table 8.2.3 includes general estimates of nutrient availability for manures and composts but these can vary widely depending on animal feed, management of grazing, the age of the manure, amount and type of bedding, and many other factors. See table 3.1 for estimates of the nitrogen content of various cover crops. **Manure applications may not be allowed by your certifier or marketer even if applied 120 days before harvest. Check with both these sources prior to making manure applications.**

Table 8.2.3 Nutrient Content of Common Animal Manures and Manure Composts

	TOTAL N	P ₂ O ₅	K ₂ O	N1 ¹	N2 ²	P ₂ O ₅	K ₂ O
	NUTRIENT CONTENT LB/TON			AVAILABLE NUTRIENTS LB/TON IN FIRST SEASON			
Dairy (with bedding)	9	4	10	6	2	3	9
Horse (with bedding)	14	4	14	6	3	3	13
Poultry (with litter)	56	45	34	45	16	36	31
Composted dairy manure	12	12	26	3	2	10	23
Composted poultry manure	17	39	23	6	5	31	21
Pelleted poultry manure ³	80	104	48	40	40	83	43
Swine (no bedding)	10	9	8	8	3	7	7
	NUTRIENT CONTENT LB/1000 GAL.			AVAILABLE NUTRIENTS LB/1000 GAL FIRST SEASON			
Swine finishing (liquid)	50	55	25	25*	20+	44	23
Dairy (liquid)	28	13	25	14*	11+	10	23

1-N1 is an estimate of the total N available for plant uptake when manure is incorporated within 12 hours of application, 2-N2 is an estimate of the total N available for plant uptake when manure is incorporated after 7 days. 3 –Pelletized poultry manure compost. (Available in New York from Kreher’s.) * injected, + incorporated.

Adapted from “Using Manure and Compost as Nutrient Sources for Fruit and Vegetable Crops” by Carl Rosen and Peter Bierman (Reference 10) and Penn State Agronomy Guide 2007-8 (Reference 28a).

Tables 8.2.4-8.2.6 lists some commonly available fertilizers, their nutrient content.

Table 8.2.4 Available Nitrogen in Organic Fertilizer

Sources	Pounds of Fertilizer/Acre to Provide X Pounds of N per Acre				
	20	40	60	80	100
Blood meal, 13% N	150	310	460	620	770
Soy meal 6% N (x 1.5)* also contains 2% P and 3% K ₂ O	500	1000	1500	2000	2500
Fish meal 9% N, also contains 6% P ₂ O ₅	220	440	670	890	1100
Alfalfa meal 2.5% N also contains 2% P and 2% K ₂ O	800	1600	2400	3200	4000
Feather meal, 15% N (x 1.5)*	200	400	600	800	1000
Chilean nitrate 16% N cannot exceed 20% of crop’s need.	125	250	375	500	625

* Application rates for some materials are multiplied to adjust for their slow to very slow release rates.

Table 8.2.5 Available Phosphorous in Organic Fertilizer

SOURCES	Pounds of Fertilizer/Acre to Provide X Pounds of P ₂ O ₅ Per Acre				
	20	40	60	80	100
Bonemeal 15% P ₂ O ₅	130	270	400	530	670
Rock Phosphate 30% total P ₂ O ₅ (x4)*	270	530	800	1100	1300
Fish meal, 6% P ₂ O ₅ (also contains 9% N)	330	670	1000	1330	1670

* Application rates for some materials are multiplied to adjust for their slow to very slow release rates.

Table 8.2.6 Available Potassium in Organic Fertilizers.

SOURCES	Pounds of Fertilizer/Acre to Provide X Pounds of K ₂ O per acre:				
	20	40	60	80	100
Sul-Po-Mag 22% K ₂ O also contains 11% Mg	90	180	270	360	450
Wood ash (dry, fine, grey) 5% K ₂ O, also raises pH	400	800	1200	1600	2000
Alfalfa meal 2% K ₂ O also contains 2.5% N	1000	2000	3000	4000	5000
Greensand or Granite dust 1% K ₂ O (x 4)*	8000	16000	24000	32000	40000
Potassium sulfate 50% K ₂ O	40	80	120	160	200

* Application rates for some materials are multiplied to adjust for their slow to very slow release rates. Tables 8.4 to 8.6 adapted by Vern Grubinger from the University of Maine soil testing lab (Link 27).

An example of how to determine nutrient needs for beans.

You will be growing an acre of snap beans. The Cornell Integrated Crop and Pest Management Guidelines suggests a total of 40 lb. N, 80 lb. P, and 60 lb. K per acre. Your soil tests show a pH of 6.5, with medium P and K levels and recommends 60 lbs P₂O₅/acre and 40 lbs K₂O/acre (see Table 8.2.2). The field you will plant has 3% organic matter which will supply approximately 60 lbs N (line 3a). Last fall you spread but did not promptly incorporate 5 tons/acre of dairy manure with bedding before planting a rye cover crop supplying ~10 lbs N (line 3d). Nutrient credits for soil organic matter, manure, and cover crop appear in Table 8.2.7.

Table 8.2.7 Bean Example: Calculating Nutrient Credits and Needs Based on Soil Sample Recommendations.

	Nitrogen (N) lbs/acre	Phosphate (P ₂ O ₅) lbs/acre	Potash (K ₂ O) lbs/acre
1. Total crop nutrient needs:	40	80	60
2. Recommendations based on soil test	Not provided	60	40
3. Credits			
a. Soil organic matter 3%	60	-	-
b. Manure – 5 ton dairy	10	15	45
c. Compost - none	0	0	0
d. Cover crop – rye	0	0	0
4. Total credits:	70	15	45
5. Additional needed (2-4) =	0	45	0

Because of the threat of seedcorn maggot and possible N tie-up from the decomposing rye cover, wait at least three weeks between plowing and planting. This would be a good opportunity to do a shallow tine weeding just before planting. The dairy manure and soil organic matter will release N and K at adequate levels in this example; P is the only nutrient that needs to be applied. Because beans are sometimes planted into cold soils consider applying the 45 lbs/acre of P in a relatively soluble form such as composted chicken manure, which contains about 100 lbs P₂O₅ per ton. Thus, banding about 900 lb at planting (2-3 inches to the side and below the furrow) would provide the P needed in this example, but would provide 40 lb of additional N, which is already higher than needed. Banding about 600 lbs of rock phosphate instead could avoid too much N, but the P would be less available in colder soils.

9. HARVESTING

All processing snap bean acreage is harvested by machine. Snap beans are processed relatively soon after harvest.

10. USING ORGANIC PESTICIDES

Given the high cost of many pesticides and the limited amount of efficacy data from replicated trials with organic products, the importance of developing an effective system of cultural practices for insect and disease management cannot be emphasized strongly enough. **Pesticides should not be relied on as a primary method of pest control.** Scouting and forecasting are important for detecting symptoms of diseases at an early stage. When conditions do warrant an application, proper choice of materials, proper timing, and excellent spray coverage are essential.

10.1 Sprayer Calibration and Application

Calibrating sprayers is especially critical when using organic pesticides since their effectiveness is sometimes limited. For this reason, they tend to require the best spraying conditions to be effective. Read the label carefully to be familiar with the unique requirements of some products, especially those with live biological organisms as their active ingredient (e.g. Contans). The active ingredients of some biological pesticides (e.g. Serenade and Sonata) are actually metabolic byproducts of the organism. Calculating nozzle discharge and travel speed are two key components required for applying an accurate pesticide dose per acre. Applying too much pesticide is illegal, can be unsafe and is costly whereas applying too little can fail to control pests or lead to pesticide resistance.

Resources

- [Cornell Integrated Crop and Pest Management Guidelines: Pesticide Information and Safety](#) (Reference 62).
- [Calibrating Backpack Sprayers](#) (Reference 63).
- [Agricultural Pocket Pesticide Calibration Guide](#) (Reference 64).
- [Knapsack Sprayers – General Guidelines for Use](#) (Reference 65)
- [Herbicide Application Using a Knapsack Sprayer](#) (Reference 66) this publication is relevant for non-herbicide applications).

10.2 Regulatory Considerations

Organic production focuses on cultural, biological, and mechanical techniques to manage pests on the farm, but in some cases organically approved pesticides, which include repellents, are a necessary option. Pesticides mentioned in this organic production guide must be registered and labeled at the federal level for use, like any other pesticide, by the Environmental Protection Agency (EPA), or meet the EPA requirements for a “minimum risk” pesticide, making it exempt from normal registration requirements as described in [FIFRA regulation 40 CFR Part 152.25\(b\)](#) (Link 67).

“Minimum risk” pesticides, also referred to as 25(b) pesticides, must meet specific criteria to achieve the “minimum risk” designation. The active ingredients of a minimum-risk pesticide must be on the list of exempted active ingredients found in the federal regulations (40 CFR 152.25). Minimum-risk pesticides must also contain inert ingredients listed on the most [current List 4A](#) published in the Federal Register (Link 68).

In addition to meeting the active and inert ingredient requirements above, a minimum-risk pesticide must also meet the following:

- Each product must bear a label identifying the name and percentage (by weight) of each active ingredient and the name of each inert ingredient.
- The product must not bear claims to either control or mitigate microorganisms that pose a threat to human health, including, but not limited to, disease-transmitting bacteria or viruses, or claim to control insects or rodents carrying specific diseases, including, but not limited to, ticks that carry Lyme disease.
- The product must not include any false or misleading labeling statements.

Besides registration with the EPA, pesticides sold and/or used in New York State must also be registered with the New York State Department of Environmental Conservation (NYS DEC). However, pesticides meeting the EPA “minimum risk” criteria described above do not require registration with the NYS DEC.

To maintain organic certification, products applied must also comply with the National Organic Program (NOP) regulations as set forth in [7 CFR Part 205, sections 600-606](#) (Link 69). The Organic Materials Review Institute (OMRI) (Link 5) is one organization that reviews and publishes products they find compliant with the NOP regulations, but other entities also make product assessments. Organic growers are not required to use only OMRI listed materials, but the list is a good starting point when searching for potential pesticides.

Finally, each farm must be certified by an accredited certifier who must approve any material applied for pest management. ALWAYS check with the certifier before applying any pest control products.

Some organic certifiers may allow "home remedies" to be used to manage pests. These materials are not labeled as pesticides, but may have properties that reduce the impact of pests on production. Examples of home remedies include the use of beer as bait to reduce slug damage in strawberries or dish detergent to reduce aphids on plants. Home remedies are

not mentioned in these guides, but in some cases, may be allowed by organic certifying agencies. Maintaining good communication with your certifying agent cannot be overemphasized in order to operate within the organic rules.

10.3 Optimizing Pesticide Effectiveness

Information on the effectiveness of a particular pesticide against a given pest can sometimes be difficult to find. Some university researchers include pesticides approved for organic production in their trials; some manufacturers provide trial results on their web sites; some farmers have conducted trials on their own. Efficacy ratings for pesticides listed in this guide were summarized from university trials and are only provided for some products. Pesticide manufacturers are not required to demonstrate efficacy to list a pest on the label. The [Resource Guide for Organic Insect and Disease Management](#) (Reference 2) provides efficacy information for many approved materials.

In general, pesticides allowed for organic production may kill a smaller percentage of the pest population, could have a shorter residual, and may be quickly broken down in the environment. Read the pesticide label carefully to determine if water pH or hardness will negatively impact the pesticide’s effectiveness. Use of a surfactant may improve organic pesticide performance. [OMRI lists adjuvants](#) on their website under *Crop Management Tools and Production Aids* (Link 5). Regular scouting and accurate pest identification are essential for effective pest management. Thresholds used for conventional production may not be useful for organic systems because of the typically lower percent mortality and shorter residual of pesticides allowed for organic production. When pesticides are needed, it is important to target the most vulnerable stages of the pest. Thoroughly cover plant surfaces, especially in the case of insecticides, since many must be ingested to be effective. The use of pheromone traps or other monitoring or prediction techniques can provide an early warning for pest problems, and help effectively focus scouting efforts.

11. DISEASE MANAGEMENT

In organic systems, cultural practices form the basis of a disease management program. Promote plant health by maintaining a biologically active, well-structured, adequately drained and aerated soil that supplies the requisite amount and balance of nutrients. Choose varieties resistant to one or more important diseases whenever possible. Plant only clean, disease-free seed and maintain the best growing conditions possible.

Rotation is an important management practice for pathogens that overwinter in crop debris. Rotating between crop families is useful for many diseases, but may not be effective for pathogens with a wide host range, such as *Sclerotinia* white mold. Rotation with a grain crop, preferably a sod that will be in place for one or more seasons, deprives disease-causing organisms of a host, and also contributes to a healthy soil structure that promotes vigorous plant growth. The same practices are effective for preventing the buildup of root damaging nematodes in the soil, but keep in mind that certain grain crops are also hosts for some nematode species. See more on crop rotation in Section 4.2: *Crop Rotation Plan*.

Other important cultural practices can be found under each individual disease listed below. Maximizing air movement and leaf drying is a common theme. Many plant diseases are favored by long periods of leaf wetness. Any practice that promotes faster leaf drying, such as orienting rows with the prevailing wind, or using a wider row or plant spacing, can slow disease development. Fields surrounded by trees or brush that tend to hold moisture after rain, fog or dew, should be avoided if possible.

Scouting fields weekly is key to early detection and evaluating control measures. The earlier a disease is detected, the more likely it can be suppressed with organic fungicides. When available, scouting protocols can be found in the sections listed below for each individual disease. While following a systematic scouting plan, keep watch for other disease

problems. Removing infected plants during scouting is possible on a small operation. Accurate identification of disease problems, especially recognizing whether they are caused by a bacterium or fungus, is essential for choosing an effective control strategy. Anticipate which diseases are likely to be problems that could affect yield and be ready to take control action as soon as symptoms are seen. Allowing pathogen populations to build can quickly lead to a situation where there are few or no options for control.

All currently available fungicides allowed for organic production are protectants meaning they must be present on the plant surface before disease inoculum arrives to effectively prevent infection. They have no activity on pathogens once they are inside the plant. A few fungicides induce plant resistance and must be applied several days in advance of infection to be effective. Biological products must be handled carefully to keep the microbes alive. Follow label instructions carefully to achieve the best results.

Contact your local cooperative extension office to see if newsletters and pest management updates are available for your region. For example, the Cornell Cooperative Extension Regional Vegetable Program in Western New York offers subscriptions to *Pestminder*, a report that gives timely information regarding crop development, pest activity and control, and *VegEdge*, a monthly newsletter with articles on pest management. In the Albany area, information is available through the weekly *Vegetable Pest Status Report*. On Long Island, see the *Long Island Fruit and Vegetable Update*.

Organic farms must comply with all other regulations regarding pesticide applications. See Section 10. Using Organic Pesticides for details. **ALWAYS check with your organic farm certifier when planning pesticide applications.**

Resources:

[Cornell Vegetable MD Online](#) (Link 13)

[Resource Guide for Organic Insect and Disease Management](#) (Reference 2)

ORGANIC SNAP BEAN PRODUCTION

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 11.0 Pesticides Labeled for Organic Bean Disease Control

Class of Compound Product name - <i>Active ingredient</i>	Bacterial Diseases			Bean Rust	Botrytis Grey Mold	Pod-Flecking Complex	Root Rot & Damping Off	Sclerotinia White Mold
	Bacterial Brown Spot	Common Bacterial Blight	Halo blight					
BIOLOGICALS								
ActinoGrow - <i>Streptomyces lydicus</i>					X	X	X	X
Actino-Iron - <i>Streptomyces lydicus</i>							X	X
Actinovate AG - <i>Streptomyces lydicus</i>					X	X		X
Contans WG - <i>Coniothyrium minitans</i>								2
Mycostop - <i>Streptomyces griseoviridis</i>							X	
Mycostop Mix - <i>Streptomyces griseoviridis</i>							X	
Serenade ASO - <i>Bacillus subtilis</i>				X				1
Serenade MAX - <i>Bacillus subtilis</i>				X				X
T-22 – <i>Trichoderma harzianum</i>							X	
BOTANICALS								
Sporan EC - <i>Herbal oils</i>	X	X	X	X	X			
Trilogy - <i>Neem extract</i>				X	1	X		
COPPER								
Basic Copper 53 - <i>Copper sulfate</i>	2	2	2					
Champ WG – <i>Copper hydroxide</i>	X	X	X					
Cueva Fungicide Concentrate – <i>copper octanoate</i>	X	X	X		X			X
Nu Cop 50 WP - <i>Copper hydroxide</i>	2	2	2					
SULFUR								
MicroSulf - <i>sulfur</i>	X			X				
Microthiol Disperss - <i>Sulfur</i>				X				
OIL								
Organic JMS Stylet Oil - <i>Paraffinic oil</i>				X				
OTHER								
Heads Up Plant Protectant – <i>Chenopodium quinoa</i>							X	

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, X- Labeled but not reviewed or research not available.

11.1 Sclerotinia White Mold, *Sclerotinia sclerotiorum*

Time for concern: From open blossoms through the end of harvest

Key characteristics: The fungus will initially attack bean blossoms, as blossoms are a readily available source of food. Symptoms appear as white, fluffy cottony growth on blossoms, stems and pods. As the fungus grows, mounds of white mycelium harden and darken. These dark, black structures become sclerotia that enable the fungus to overwinter. Bean blossoms are an excellent source of nutrients for the fungus. Therefore control measures must be initiated at bloom. See Link 31. See Cornell [photo](#), and [bulletin](#) (Links 32-33) and learn more at the Dillard Lab Vegetable Pathology website at nysaes.cornell.edu/pp/faculty/dillard/main.html.

Management Option	Recommendation for White Mold
Scouting/thresholds	Scout field prior to harvest to determine the need for treatment with Contans WG after harvest to reduce overwintering inoculum. Keep an accurate history of white mold incidence and severity in all fields.
Coverage	The best coverage can be obtained by using a minimum of 50 gallons per acre and high pressure (100 to 200 psi). Thoroughly cover initials, buds, and blossoms.
Resistant varieties	No resistant varieties are available, however plant architecture influences disease development. Select varieties with open canopies that hold pods high off the ground.
Crop rotation	If there is a field history of white mold, beans should not be preceded by a bean, tomato, potato, lettuce, or crucifer crop for several years. Grains and corn are good rotation crops.
Site selection	Avoid planting in shaded areas and in small fields surrounded by trees; do not plant in fields that drain poorly or have a history of severe white mold.
Planting	Plant rows in an east-west direction and use wide row spacing, 36 inches, to promote drying of the soil and reduce moisture in the plant canopy.
Fertilization	Avoid over fertilization.
Postharvest	Incorporate crop debris immediately following harvest to allow soil microorganisms the opportunity to feed on the survival structures called sclerotia or degrade disease organisms/overwintering structures.
Seed selection/treatment	This is not currently a viable management options.
Note(s)	White mold tends to develop in dense plant canopies. The disease tends to be worse in fields where there is poor weed management, leaves have mechanical damage or pesticide injury, and where dead leaves are on the ground. The fungus can grow on dead and living material. White mold tends to develop when wet weather is persistent.

ORGANIC SNAP BEAN PRODUCTION

Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 11.1 Pesticides Labeled for Management of Sclerotinia White Mold					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
BIOLOGICALS					
ActinoGrow (<i>Streptomyces lydicus</i> (WYEC 108))	1-6oz/A foliar 1-12oz/A drench 2-18oz/100lb seed*	0	1 or when dry	?	*can be applied to seed in a slurry or dry in planter box.
Actino-Iron (<i>Streptomyces lydicus</i> (WYEC 108))	10-15 lb	-	4	?	Water in after application
Actinovate AG (<i>Streptomyces lydicus</i> WYEC 108)	3-12 oz	0	1*	?	*1 hour re-entry or until solution is dry. Reapply every 7-14 days. Use a spreader-sticker. Use as foliar spray or soil drench.
Contans (<i>Coniothyrium minitans</i>)	2-4 lb	-	4	2	Effective in 4/7 trials. Apply Contans to a Sclerotinia-infected crop immediately following harvest at 1 lb/A and incorporate the debris into the soil or apply at 2 lb/acre to a planted crop right after planting followed by shallow incorporation (or irrigate) to about a 1 to 2 inch depth. Do not turn the soil profile after application of Contans to avoid bringing untreated soil containing viable sclerotia near the surface. The seller recommends applying Contans for at least 3 to 4 years to reduce soil levels, or every year a susceptible crop is grown in that field. Enhance storage life by keeping product in the refrigerator or freezer.
Serenade ASO (<i>Bacillus subtilis</i>)	2-6 quarts	0	4	1	Effective in 1/1 trial.
Serenade MAX (<i>Bacillus subtilis</i>)	1-3 lbs	0	4	4	
COPPER					
Cueva Fungicide Concentrate (<i>copper octanoate</i>)	0.5-2.0 gal/100gal water	Up to day of harvest	4	?	Note that mixed material is applied at 50-100 gallons of diluted spray per acre.

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.
PHI = pre-harvest interval, REI = restricted-entry interval.

11.2 Botrytis Gray Mold, *Botrytis cinerea*

Time for concern: From open blossoms through the end of harvest

ORGANIC SNAP BEAN PRODUCTION

Key characteristics: Gray mold develops in dense plant canopies when the weather is warm and moist. Large necrotic lesions produce gray spores on the pods, leaves and stems. This disease is worse in fields where leaves have mechanical damage or other injury. See Cornell [photo](#) (Links 34) and learn more at the Dillard Lab Vegetable Pathology website at nysaes.cornell.edu/pp/faculty/dillard/main.html.

Management Option	Recommendations for Botrytis Gray Mold
Scouting/thresholds	Botrytis cinerea can infect many species of plants. A reliable forecasting system has not been developed for gray mold on beans. However, here are a few helpful hints on the development of gray mold. Gray mold tends to develop in dense plant canopies. The disease tends to be worse in fields where leaves have mechanical damage and where dead leaves are on the ground. The fungus grows and produces spores on dead and living material. These spores will subsequently infect bean pods. Gray mold tends to develop when the weather is warm and moist. Begin scouting when the first buds are showing. Record the occurrence and severity of gray mold. See Reference 9.
Coverage	The best coverage can be obtained by using a minimum of 50 gallons per acre and high pressure (100 to 200 psi).
Resistant varieties	No resistant varieties are available.
Crop rotation	Regular crop rotation is recommended. Grains and corn are good rotation crops.
Site selection	Avoid planting in shaded areas and in small fields surrounded by trees; do not plant in fields that drain poorly.
Planting	Plant rows in an east-west direction, and use wide row spacing (36 inches) to promote drying of the soil and reduce moisture in the plant canopy.
Fertilization	Avoid over-fertilization with nitrogen.
Postharvest	Incorporate debris immediately after harvest to hasten decomposition of the material.
Seed selection/treatment	This is not currently a viable management options.
Note(s)	Avoid mechanical damage to leaves.

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 11.2 Pesticides Labeled for Management of Botrytis Gray Mold					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
BIOLOGICALS					
ActinoGrow (<i>Streptomyces lydicus</i> (WYEC 108)	1-6oz/A foliar	0	1 or when dry	?	
Actinovate AG (<i>Streptomyces lydicus</i>)	3-12 oz (soil treatment)	0	1*	?	*1 hour re-entry or until solution is dry. Foliar spray must be reapplied every 7-14 days using a spreader sticker. Use as soil drench.
BOTANICAL					

ORGANIC SNAP BEAN PRODUCTION

Table 11.2 Pesticides Labeled for Management of Botrytis Gray Mold					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
Sporan EC (Herbal oils)	1.0-3.0 pts in a minimum of 25 gal water/A	0	0	?	
Trilogy (Neem extract)	0.5-1.0% in 25-100 gal water/A	None listed	4	1	Effective in 1/1 trial. Apply sufficient water to achieve complete coverage of foliage. Repeat applications every 7-14 days. Maximum labeled use of 2 gal/acre/application.
COPPER					
Cueva Fungicide Concentrate (copper octanoate)	0.5-2.0 gal/100gal water	Up to day of harvest	4	?	Note that mixed material is applied at 50-100 gallons of diluted spray per acre.

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.
PHI = pre-harvest interval, REI = restricted-entry interval.

11.3 Pod-flecking complex (PFC), *Alternaria alternata* and *Plectosporium tabacinum*

Time for concern: From pod fill through the end of harvest

Key characteristics: Pod-flecking complex is sometimes referred to by consultants and producers as russet, seam rust, spots, or rusty or spotty beans. PFC symptoms on pods intensify with pod maturity, and are most prevalent mid- to late August following periods of prolonged rainfall or rainfall of high intensity. Symptoms on pods include tan, orange, or black discolorations in the suture and/or small dark superficial specks, flecks, or spots (sometimes sunken) on the pod surfaces.

See <http://www.nysaes.cornell.edu/pp/faculty/dillard/beans.html>.

Management Options	Recommendations for Pod-Flecking Complex
Scouting/thresholds	Pod-flecking complex can infect many species of plants. A reliable forecasting system has not been developed for PFC on beans. However, here are a few helpful hints on the development of PFC. Pod-flecking complex symptoms on pods intensify with pod maturity, and are most prevalent mid- to late August following periods of prolonged rainfall or rainfall of high intensity. Only the pods are affected. Symptoms include tan, orange, or black discolorations in the suture and/or small dark superficial specks, flecks, or spots (sometimes sunken) on the pod surfaces. Begin scouting at pod fill. Record the occurrence and severity of PFC. See Reference 18.
Coverage	The best coverage can be obtained by using a minimum of 50 gallons per acre and high pressure (100 to 200 psi).
Resistant varieties	No resistant varieties are available.
Crop rotation	Regular crop rotation is recommended to improve plant health.
Site selection	Avoid planting in shaded areas and in small fields surrounded by trees; do not plant in fields that drain poorly.
Planting	Plant rows in an east-west direction, and use wide row spacing (36 inches) to promote drying of the soil and reduce moisture in the plant canopy.
Fertilization	Avoid over-fertilization with nitrogen.
Harvest / Postharvest	To mitigate disease, harvest at or near peak maturity and avoid harvest delays that would result in overripe pods. Don't store infected beans.

ORGANIC SNAP BEAN PRODUCTION

Seed selection/treatment	Incorporate debris immediately after harvest to hasten decomposition of the material. This is not currently a viable management options.
---------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 11.3 Pesticides Labeled for Management of Pod-Flecking Complex					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
BIOLOGICALS					
ActinoGrow (<i>Streptomyces lydicus</i> WYEC 108)	1-6oz/A foliar	0	1 or when dry	?	Labeled for alternaria only.
Actinovate AG (<i>Streptomyces lydicus</i> WYEC 108)	3-12 oz (foliar spray)	0	1*	?	Labeled for alternaria only. *1 hour re-entry or until solution is dry. Reapply every 7-14 days. The Label recommends use of a spreader-sticker for foliar sprays.
BOTANICAL					
Trilogy (<i>Neem extract</i>)	0.5-1.0% in 25-100 gal water/A	None listed	4	?	Maximum labeled use of 2 gal/acre/application.

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.

PHI = pre-harvest interval, REI = restricted-entry interval.

11.4 Root Rot and Damping-Off.

There are four pathogenic fungi, Pythium, Rhizoctonia, Thielaviopsis, and Fusarium that are often found attacking bean roots and causing yield losses.

Time for concern: Planting through end of bloom stage. Long term planning is required for sustainable management.

Key characteristics: Pythium is seen early in the season when it causes rotting of seeds and damping-off of young seedlings. In older plants, it causes reduction and discoloration of the root system. Pythium can also attack pods, causing lesions that generally develop at the tip of the pod and expand toward the stem. See Cornell [pythium photo](#) (Link 36). Rhizoctonia is also capable of causing seed decay and damping-off diseases on seedlings. See Cornell [rhizoctonia photo](#) (Link 37). On older plants it produces reddish brown, sunken lesions on the stem and tap root, and is generally favored by warm soil conditions. In addition, the sexual stage of Rhizoctonia may also be detected on the stem, petioles, and pods as a thin, whitish, compact growth. Thielaviopsis is often referred to as black root rot because the initial elongated lesions and later large infected areas on the stems and roots are dark brown to charcoal. This disease is favored by hot, wet conditions. Fusarium rot causes longitudinal, brick red lesions on the stem and tap root and is very common. See References 9 and Link 31, and Cornell [bulletin](#) (Link 35) for more photos and information.

Management Option	Recommendations for Root Rot and Damping-Off
Scouting/thresholds	Record the occurrence, type, and severity of root rot. No thresholds are available.

ORGANIC SNAP BEAN PRODUCTION

Resistant varieties	All commercially acceptable varieties are susceptible, but a number yield better under severe root rot.
Crop rotation	Rotate away from vegetables. One or two years with a grain crop such as barley, oats, rye, wheat, or corn will prevent severe root rot development when practiced on relatively clean fields; longer rotations are necessary in heavily infested fields. Avoid planting legumes as rotational crops or cover crops in heavily infested areas.
Site selection	Choose healthy and well-drained soils. A soil-indexing procedure that differentiates relatively pathogen-free fields from those with severe root rot problems is available. See the Cornell Soil Health Assessment Manual (Link 14) for directions to do this yourself or to send a sample to Cornell for testing.
Seed selection/treatment	Select vigorous, disease-free seeds.
Soil treatment	Breaking hard pans, plowing deep, or ripping and planting on raised ridges or beds will reduce damage from root rot diseases.
Cover crop	Barley, rye grain, rye grass, wheat, oats, and other grain crops left on surface or plowed under as green manures or dry residue in the spring are beneficial. If incorporated as green manures, allow 2 weeks or more for decomposition prior to planting. Sudangrass or sorghum sudangrass hybrids can also be used as green manures. In wet years, using green manures may increase slug damage and affect stand establishment.
Planting	Avoid planting in heavily infested fields, but if there is no choice, plant shallow and late. Plantings exhibiting symptoms of severe root rot damage will benefit from a shallow cultivation not too close to the stems. Also, covering the lower stem tissues with soil will promote further root formation and reduce root rot damage. However, the latter should be done on an emergency basis, as this practice has been observed to increase foliar infections with <i>Rhizoctonia</i> in wet seasons.
Postharvest	Crop debris should be plowed down to initiate decomposition, if tillage system in use permits.

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 11.4 Pesticides Labeled for Management of Root Rot and Damping-Off					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
BIOLOGICAL					
ActinoGrow (<i>Streptomyces lydicus</i> (WYEC 108))	1-12oz/A drench 2-18oz/100lb seed*	0	1 or when dry	?	*can be applied to seed in a slurry or dry in planter box.
Actino-Iron (<i>Streptomyces lydicus</i> (WYEC 108))	10-15 lb	-	4	?	Water in after application

ORGANIC SNAP BEAN PRODUCTION

Table 11.4 Pesticides Labeled for Management of Root Rot and Damping-Off					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
Mycostop Mix (<i>Streptomyces griseoviridis</i> Strain K61)	5-8 oz/100 lbs of seed as seed treatment 7.6-30 oz/A as soil spray or drench 0.5-1 lb/ treated acre as band, in-furrow or side dress.	0	4	?	Labeled only for control of root rots and damping off caused by <i>Fusarium</i> spp. Use at planting; no pre-harvest interval noted. Irrigate within 6 hours after soil spray or drench with enough water to move Mycostop into the root zone. Lightly incorporate furrow or band applications.
Mycostop (<i>Streptomyces griseoviridis</i> Strain K61)	8 oz/cwt seed as seed treatment 15-30 oz/a as soil spray or drench	-	4	?	Irrigate within 6 hours after soil spray or drench with enough water to move Mycostop into the root zone.
T-22 (<i>Trichoderma harzianum</i>)	1.5-3 oz/cwt seed	-	0	?	Use as a hopper box seed treatment
OTHER					
Heads Up Plant Protectant (<i>Chenopodium quinoa extract</i>)	.13 oz/gal water (seed treatment)	-	12	?	Use 34 ounces of mixed product to treat 360 pounds of seed. Thoroughly coat seed surface with solution. Use immediately after mixing.

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.
PHI = pre-harvest interval, REI = restricted-entry interval.

11.5 Bacterial Diseases

Bacterial brown spot (Pseudomonas syringae pv. syringae); Common bacterial blight, (Xanthomonas campestris pv. phaseoli); and Halo blight, (Pseudomonas syringae pv. phaseolicola)

Time for concern: From seeding stage through the end of harvest

Key characteristics: Brown spot - small, brown spots, 3/25 to 8/25 inch in diameter, often with a narrow, diffuse, pale margin, appear on leaves. Sunken, brown spots can form on the pods. Common bacterial blight - light brown lesions of irregular shape with distinct, bright yellow margins, 2/5 inch long, appear on leaves. Spots form on the pods and enlarge into reddish brown lesions. In humid weather, yellow bacteria may be present on the lesions. Halo blight - small, water-soaked spots on the undersides of leaves develop into numerous, small, reddish brown lesions with pale to yellow margins or halos. Pod symptoms are similar to those of common blight. See Cornell [fact sheet](#) (Link 38).

Management Option	Recommendations for Bacterial Diseases
Scouting/thresholds	Fields should be scouted at least twice between midseason and harvest. Record the occurrence and severity of the bacterial blights.
Resistant varieties	Some varieties have tolerance and/or resistance to one or more of these diseases. Depending on the variety, brown spot development on leaves may or may not result in significant damage to pods. Wisconsin growers report that 'Hystyle' shows resistance to brown spot.
Crop rotation	In the case of halo blight, rotate away from fields where this disease has occurred for a minimum of three years. Use a two-year minimum rotation for bacterial brown spot or common bacterial blight. <i>Xanthomonas campestris</i> also harbors in weeds within the mustard family.
Site selection	Avoid planting snap bean fields near dry bean fields.

ORGANIC SNAP BEAN PRODUCTION

Management Option	Recommendations for Bacterial Diseases
Seed selection/treatment	Plant only western-grown, certified seed.
Postharvest	Crop debris should be destroyed as soon as possible to remove this source of disease for future plantings and to initiate decomposition.
Sanitation	Equipment used in fields with bacterial diseases should be thoroughly cleaned before being moved to disease-free fields. To reduce the spread of bacteria on equipment or in spray water, avoid making pesticide applications or cultivating when the leaves are wet. If possible, plow under bean stubble immediately after harvest.

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)) (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 11.5 Pesticides Labeled for Management of Bacterial Diseases					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
BOTANICAL					
Sporan (Herbal oils)	1.0-3.0 pts in a minimum of 25 gal water/A	0	0	?	Check label for specific uses for bacterial diseases.
COPPER COMPOUNDS					
Basic Copper 53 (Copper Sulfate)	2 – 4 lbs	Up to day of harvest	24	2	Control has been inconsistent*. If possible time applications so that 12 hours of dry weather follow application.
Champ WG (Copper hydroxide)	1-3 lbs	-	24	?	
Cueva Fungicide Concentrate (copper octanoate)	0.5-2.0 gal/100gal water	Up to day of harvest	4	?	Note that mixed material is applied at 50-100 gallons of diluted spray per acre.
Nu Cop 50WP (Copper hydroxide)	1.5-3 lbs	1	24	2	Control has been inconsistent*.
SULFUR					
MicroSulf (Sulfur)	7 lbs	-	24	?	Consult processor before using sulfur.

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.

PHI = pre-harvest interval, REI = restricted-entry interval.

* From Cornell Integrated Crop and Pest Management Guidelines

11.6 Virus Diseases

Virus diseases of bean are spread by infected seed or by aphids. However, controlling aphids is not effective for reducing these viruses. Virus expression is lower when the crop has adequate water. Irrigating in dry years may help mitigate impacts of virus infection. Record the occurrence and severity of any viruses present. See Cornell [photos](#) (Link 40), [fact sheet](#) (Link 39), and a [list of weed and crop hosts](#) (Links 41). Information about aphid vector activity and spread of CMV in snap bean fields can be found in References 3 and 6. Cornell research on developing CMV-resistant snap bean varieties is on going.

Disease/Symptoms	Spread by	Time for concern	Resistant Varieties	Notes
<p>Cucumber Mosaic Virus (CMV)</p> <p>Leaf curl, green mottle, blistering, and a rugged zipper-like appearance along the main veins involving only a few leaves. Infected plants may also not express symptoms.</p>	<p>Soybean aphid, yellow clover aphid, pea aphid, corn leaf aphid; rarely seed transmitted</p>	<p>Primarily in plantings from late June through late July.</p>	<p>None available</p>	<p>Use certified, disease-free seed. CMV does not persist in plant debris, in the soil, or on equipment. For CMV and the other aphid-transmitted viruses listed below, there will be less risk of yield loss if fields are planted early (mid May to late June) than if planted after this period.</p>
<p>Bean Common Mosaic Virus (BCMV)</p> <p>Symptoms include a green mosaic and downward cupping along the main vein of each leaflet. Green vein banding, blistering, and malformation are common in leaves of the same plant. Plants are small, and pods may be mottled and malformed. Symptoms are persistent.</p>	<p>Seed, bean aphid, cowpea aphid, pea aphid, potato aphid, green peach aphid</p>	<p>Seeding through harvest</p>	<p>Most varieties carry the I-gene for resistance to BCMV.</p>	<p>Use certified, disease-free seed.</p>
<p>Bean Yellow Mosaic Virus (BYMV)</p> <p>Leaf mosaic formed by contrasting yellow or green mosaic areas. Pods generally are not affected, but the number of seeds per pod may be reduced.</p>	<p>Bean aphid, cowpea aphid, pea aphid, potato aphid, and green peach aphid.</p>	<p>June through harvest</p>	<p>None available</p>	<p>BYMV does not persist in plant debris, in the soil, or on equipment.</p>
<p>Clover Yellow Vein Virus (CYVV)</p> <p>In addition to deforming pods, this virus also causes a prominent yellow mosaic, malformation, and reduction in plant size.</p>	<p>Aphids</p>	<p>June through harvest</p>	<p>None available</p>	

ORGANIC SNAP BEAN PRODUCTION

11.7 Bean Rust, *Uromyces appendiculatus*

Time for concern: Early in the growing season

Key characteristics: Regular occurrence of dew favors infection and development of severe epidemics. Bean rust is characterized by reddish brown, circular pustules on leaves or pods. See Cornell [photo](#) and University of Connecticut [fact sheet](#) (Links 42 and 43). Bean rust is rarely seen in New York. However, growers commonly use the name “rust” to describe a diffuse light brown discoloration that sometimes occurs on the pods. See section 10.3 on Pod-flecking complex (Reference 18). Also see the Dillard Lab Vegetable Pathology Website at <http://www.nysaes.cornell.edu/pp/faculty/dillard/main.html>.

Management Option	Recommendations for Bean Rust
Scouting/thresholds	Record the occurrence and severity of bean rust. No thresholds are available.
Resistant varieties	No resistant varieties are available.
Crop rotation	A minimum one-year rotation is recommended.
Site selection	Avoid areas with poor air and soil-moisture drainage.
Planting	Wider row spacing reduces leaf wetness and may slow epidemic development.
Postharvest	Incorporate infested debris immediately after harvest to hasten decomposition of the material.
Sanitation	Avoid walking through the crop when the leaves are wet.

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide’s effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 11.7 Labeled Products for Management of Bean Rust

Class of Compound Product (Active ingredient)	Rate/Acre	PHI (days)	REI (hours)	Efficacy	Comments
BIOLOGICALS					
Serenade ASO (<i>Bacillus subtilis</i>)	2-6 qts	0	4	?	
Serenade MAX (<i>Bacillus subtilis</i>)	1-3 lbs	0	4	?	
BOTANICALS					
Sporan EC (Herbal Oils)	1-3 qts	0	0	?	
SULFUR					
MicroSulf (Sulfur)	7 lbs	0	24	?	Consult processor before using sulfur)
Microthiol Disperss (Sulfur)	3-10 lbs	-	24	?	Do not apply within 2 weeks of an oil application nor at temperatures over 90 degrees.
OTHER					
Organic JMS Stylet Oil (Paraffinic oil)	3-6 qt/100 gal	0	4	?	See label for restrictions when combined with sulfur.
Trilogy (Neem extract)	0.5-1.0% in 25-100 gal water/A	None listed	4	?	Maximum labeled use of 2 gal/acre/application.

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.
PHI = pre-harvest interval, REI = restricted-entry interval.

12. Root-Lesion Nematode, *Pratylenchus penetrans*

Time of concern: Before planting. Long term planning is required for sustainable management

Key characteristics: The root lesion nematode typically does not cause characteristic symptoms on snap bean roots. However, severely infected plants may show general chlorosis and stunting as well as a reduced root system. Infection by this nematode may predispose plants to infection and damage by other soil borne pathogens. See Cornell [sampling for root lesion nematode](#) for more information and photos of damage (Link 46).

Management Option	Recommendations for Root-Lesion Nematode
Scouting/thresholds	Use a soil bioassay with soybean to assess soil root-lesion nematode infestation levels. Or, submit the soil sample(s) for nematode analysis at a public or private nematology lab (Link 44). See Section 4: Field Selection for more information as well as the following Cornell publications for instructions: "How to" instructions for soil sampling for nematode bioassays (Link 45). "How to" instructions for farmers to conduct a field test for root lesion nematode using soybean (Link 46).
Resistant Varieties	No resistant snap bean varieties are available.
Crop Rotation	Root-lesion nematode has over 400 hosts including many vegetable and grain crops that are planted in rotation with snap bean thus making it difficult to manage lesion nematode strictly using crop rotation once populations have reached damaging levels. Depending on the size of the infested site, marigold varieties such as 'Polynema' and 'Nemagone' are very effective at reducing nematode populations, where marigold can be established successfully.
Site Selection	Assay soil for nematode infestation, if needed.
Biofumigant Cover Crops	Cover crops with a biofumigant effect, used as green manure, may be used for managing root-lesion nematode. It is important to note that many biofumigant crops including Sudangrass, white mustard, and rapeseed are hosts to root-lesion nematode and will increase the population until they are incorporated into the soil as a green manure, at which point their decomposition products are toxic to nematodes. Research has suggested that Sudangrass hybrid 'Trudan 8' can be used effectively as a biofumigant to reduce root-lesion nematode populations. Cover crops such as forage pearl millet 'CFPM 101' and 'Tifgrain 102', rapeseed 'Dwarf Essex', and ryegrass 'Pennant' are poor hosts, and thus will limit the build-up or reduce root-lesion nematode populations when used as a "standard" cover crop.
Sanitation	Avoid moving soil from infested fields to un-infested fields via equipment and vehicles, etc. Also limit/avoid surface run-off from infested fields.
Weed control	Many common weed species including lambsquarters, redroot pigweed, common purslane, common ragweed, common dandelion and wild mustard are also hosts therefore effective weed management is also important.

13. INSECT MANAGEMENT

Effective insect management relies on accurate identification of pests and beneficial insects, an understanding of their biology and life cycle, knowledge of economically important levels of pest damage, and a familiarity with the effectiveness of allowable control practices, in other words, Integrated Pest Management (IPM).

Regular scouting and accurate pest identification are essential for effective insect management. Thresholds used for conventional production may not be useful for organic systems because of the typically lower percent mortality and shorter residual of control products allowed for organic production. The use of pheromone traps or other monitoring and prediction techniques can provide an early warning for pest problems, and help effectively focus scouting efforts.

The contribution of crop rotation as an insect management strategy is highly dependent on the mobility of the pest. Crop rotation tends to make a greater impact on reducing pest populations if the pest has limited mobility. In cases where the insects are highly mobile, leaving a greater distance between past and present plantings is better.

Natural Enemies

Learn to identify naturally occurring beneficial insects, and attract and conserve them in your fields by providing a wide variety of flowering plants in or near the field and by avoiding use of broad-spectrum insecticides during periods when natural enemies are present. In most cases, a variety of natural enemies are present in the field, each helping to reduce pest populations. The additive effects of multiple species of natural enemies, attacking different host stages, is more likely to make an important contribution to reducing pest

populations than individual natural enemy species operating alone. Natural enemies need a reason to be present in the field, either a substantial pest population, alternative hosts, or a source of pollen or nectar, and may not respond to a buildup of pests quickly enough to keep pest populations below damaging levels. Releasing insectary-reared beneficial organisms into the crop early in the pest outbreak may help control some pests but sometimes these biocontrol agents simply leave the area. For more information, see Cornell's *Natural Enemies of Vegetable Insect Pests* (Reference 69) and [A Guide to Natural Enemies in North America](#) (Link 47).

Regulatory

Organic farms must comply with all other regulations regarding pesticide applications. See Section 10 for details. ALWAYS check with your organic farm certifier when planning pesticide applications.

Efficacy

In general, insecticides allowed for organic production kill a smaller percentage of the pest population and have a shorter residual than non-organic insecticides. University –based efficacy testing is not available for many organic pesticides. See Section 10.3 for more information on application techniques that can optimize effectiveness.

Resources:

[Natural Enemies of Vegetable Insect Pests](#) (Reference 5).
[Biological Control: A Guide to Natural Enemies in North America](#) (Link 47).
[Resource Guide for Organic Insect and Disease Management](#) (Reference 2).

ORGANIC SNAP BEAN PRODUCTION

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide’s effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 13.0 Pesticides Labeled for Organic Bean Insect Control

	European Corn Borer	Mexican Bean Beetle	Potato Leafhopper	Seedcorn Maggot	Two-Spotted Mite	Tarnished Plant Bug
BOTANICAL						
Aza-Direct (<i>Azadirachtin</i>)	X	X	X			
AzaGuard (<i>Azadirachtin</i>)		X	X			
Azahar (<i>Azadirachtin</i>)		X	X		X	X
AzaMax (<i>Azadirachtin</i>)		X	X		X	X
Neemazad 1% EC (<i>Azadirachtin</i>)		X	X			
Neemix (<i>Azadirachtin</i>)		X	X			
PyGanic Crop Protection 1.4EC (<i>Pyrethrin</i>)	X	X	X			
PyGanic Crop Protection 5.0 EC (<i>Pyrethrin</i>)		X	X			
Safer Brand #567 (<i>Pyrethrin & Potassium salts of fatty acids</i>)		X	X		X	X
BIOLOGICALS						
Deliver (<i>Bacillus thuringiensis sp. kurstaki</i>)	X					
Entrust (<i>Spinosad</i>)	X					
Javelin (<i>Bacillus thuringiensis sp. kurstaki</i>)	X					
OILS						
Glacial Spray Fluid		X	X		X	
Golden Pest Spray Oil (<i>Soybean oil</i>)		X	X		X	X
Organic JMS Stylet Oil (<i>Paraffinic oil</i>)			X		X	
Saf-T-Side (<i>Petroleum oil</i>)		X	X		X	
SuffOil-X (<i>Petroleum oil</i>)		X	X		X	
OTHER						
SucraShield (<i>Sucrose octanoate esters</i>)			X		X	
Trilogy (<i>Neem extract</i>)					X	

X-labeled for use in NYS and also listed on the Organic Materials Review Institute

13.1 Mexican Bean Beetle

Time for concern: June through September

Key characteristics: Adults are 1/4-1/3 inch long, convex, and oval /in form. They vary in color from yellow when newly emerged to a coppery brown when mature. Each wing has eight black dots in three rows across the back. Eggs are orange to yellow in color and deposited in groups of 40 to 50 on the underside of leaves. Larvae are yellow, and the bodies are covered with six rows of long, black-tipped spines. Feeding by adults and larvae results in the skeletonizing of leaves. See Cornell [fact sheet](#) and [photo of damage](#) (Links 48, 49).

Management Option	Recommendations for Mexican Bean Beetle
Scouting/thresholds	Overwintering adults move into fields and feed for 1-2 weeks and then lay their eggs. Monitor fields by scouting for adults, eggs, and larvae. Because populations vary within a field and between fields in an area, it is difficult to determine when population numbers present a threat.
Natural Enemies	Natural enemies such as parasitic flies, wasps, and predators help to control Mexican bean beetle populations. An imported parasitoid, <i>Pebibobus foveolatus</i> , can be important for control. The parasitoid does not overwinter successfully, so it must be reared and released each year. Use Reference 5 or see Cornell Biological Control: A Guide to Natural Enemies in North America for identification of natural enemies (Link 47).
Trap Crop	A small strip of early-planted beans in the previous year’s snap bean field could be used as a trap crop to attract overwintered beetles where they can be managed with foliar applications of products listed below or tilled under after adults have completed oviposition.
Resistant Varieties	No resistant varieties are available.
Planting Date	Avoiding early plantings can reduce damage.
Cultural	Avoid planting sequential crops adjacent to each other. Rotating fields as far away as possible from the previous season’s fields should help to reduce populations.
Postharvest	Soon after harvest, plow under the infested crop to prevent immature beetles from completing development on the foliage and to destroy potential overwintering beetles.

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide’s effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 13.1 Pesticides Labeled for Management of Mexican Bean Beetle					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
BOTANICAL					
Aza-Direct (Azadirachtin)	1-2 pts	0	4	1	Azadirachtin-based products effective in 4/4 trials. Aza-Direct may be tank mixed at rates as low as 4 oz/A. Maximum rate is 3.5 pt/A for heavy pest infestations.
AzaGuard (Azadirachtin)	8 oz	0	4	?	Apply with OMRI approved spray oil.
Azahar (Azadirachtin)	12-41 fl oz	0	4	?	

Table 13.1 Pesticides Labeled for Management of Mexican Bean Beetle

Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
AzaMax (Azadirachtin)	1.33 fl oz/1000 ft ²	0	4	?	
Neemazad 1% EC (Azadirachtin)	18-72 fl oz	None listed	4	?	Product works on insect larvae and nymphs.
Neemix (Azadirachtin)	4-16 oz	-	12	1	Neemix controls larvae only; apply early and often for best control.
OILS					
Glacial Spray Fluid (Mineral oil)	0.75-1 gal/100gal	Up to day of harvest	4	?	Only for use against larvae. See label for specific application volumes
Golden Pest Spray Oil (Soybean oil)	2 gal	-	4	?	Only for use against larvae.
Saf-T-Side (Petroleum oil)	1-2 gal/100 gal water	Up to day of harvest	4	?	Only for use against larvae.
SuffOil-X (Petroleum oil)	1-2 gal/100 gal water	Up to day of harvest	4	?	Only for use against larvae. Do not mix with sulfur products.
COMBINATIONS					
Pyganic Crop Protection 1.4 EC OR Pyganic Crop Protection 5.0 EC (Pyrethrins) plus Neemix 4.5 (Azadirachtin)	32oz 10oz 8 oz	0 0 -	12 12 12	1 1 1	Pyganic effective in 3/3 trials. Pyganic in combination with Neemix, provided even better protection than Pyganic alone in 1/1 trial.
Pyganic Crop Protection 1.4 EC OR Pyganic Crop Protection 5.0 EC (Pyrethrins) plus Nu Film P (Spreader Sticker)	32 oz 10oz 4-6 oz	0 0	12 12 12	1 1 1	Pyganic in combination with Nufilm provided better protection than Pyganic alone in 1/1 trial.
Safer Brand #567 Pyrethrin & Insecticidal Soap Concentrate II (Pyrethrin & potassium salts of fatty acids)	1 gal of mixed spray/700 ft ² of plant surface area	Until spray has dried	12	?	See label for specific mixing instructions.

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.

PHI = pre-harvest interval, REI = restricted-entry interval.

13.2 Potato Leafhopper (PLH), *Empoasca fabae*

Time for concern: Early June through pre-bloom

Key characteristics: The adult is wedge-shaped, iridescent green in color, and 1/8 inch long. The body is widest at the head. See Cornell [fact sheet](#) for photo (Link 50). Eggs are laid singly on the underside of leaves. Both adults and nymphs are very active, running forwards, backwards, or sideways. The symptoms produced by feeding have been termed “hopperburn.” The first sign of hopperburn is whitening of the veins. These areas become flaccid and yellow in color, then desiccate, turn brown, and die. Leaf curling is also very common. The entire process takes four to five days. See Cornell [photo of damage](#) (Link 51) or Reference 16.

Management Option	Recommendations for Potato Leafhopper
Resistant varieties	Resistance of the varieties currently grown is not known.
Scouting/thresholds	Potato leafhoppers migrate from southern areas each year and their time of arrival varies. Check for the presence of adult potato leafhoppers by using a sweep net or by placing yellow, sticky traps near field edges. Nymphs are best sampled by visual examination of the undersides of leaves on the lower half of the plant. Bean yields are most likely to be reduced by potato leafhoppers if damage occurs before bloom. Management should occur when a threshold is met of one nymph per trifoliolate leaf or when adults exceed 100/20 sweeps. On newly emerged beans, lower densities of leafhoppers than those mentioned above may be damaging. See Reference 16.
Natural enemies	Although a variety of natural enemies of potato leafhoppers have been reported, their impact on infestations is not well known. Use Reference 5 or see Cornell guide to natural enemies (Link 47).

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide’s effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 13.2 Pesticides Labeled for Management of Potato Leafhopper					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
BOTANICAL					
Aza-Direct (Azadirachtin)	1-2 pints	0	4	1	Azadirachtin-based product was effective in 1 trial.
AzaGuard (Azadirachtin)	10 oz	0	4	?	Apply with OMRI approved spray oil.
Azahar (Azadirachtin)	10-41 fl oz	0	4	?	
AzaMax (Azadirachtin)	1.33 fl oz/1000 ft ²	0	4	?	
Neemazad 1% EC (Azadirachtin)	31-72 fl oz	Not listed	4	?	Product works on insect larvae and nymphs.
Neemix 4.5 (Azadirachtin)	7-16 oz	-	12	2	Neemix alone effective in 1/2 trials. Neemix in combination with Pyganic was effective in 1/1 trial.
Pyganic Crop Protection 1.4 EC (Pyrethrin)	16-64 oz	0	12	2	Pyganic alone effective in 2/3 trials. Two applications of Pyganic may be needed to reduce adult and nymph leafhopper populations. Pyganic in combination with Neemix effective in 1/1 trial.
Pyganic Crop Protection 5 EC (Pyrethrin)	4.8-18 oz	0	12	?	

Table 13.2 Pesticides Labeled for Management of Potato Leafhopper					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
Safer Brand #567 Pyrethrin & Insecticidal Soap Concentrate II (Pyrethrin & potassium salts of fatty acids)	1 gal of mixed spray/700 ft ² of plant surface area	-	12	?	See label for specific mixing instructions.
OIL					
Glacial Spray Fluid (Mineral oil)	0.75-1 gal/100gal	Up to day of harvest	4	?	See label for specific application volumes
Golden Pest Spray Oil (Soybean oil)	2 gal	-	4	?	
Organic JMS Stylet Oil (Paraffinic oil)	3-6 qt/100 gal	0	4	?	See label for restrictions when combined with sulfur.
Saf-T-Side (Petroleum oil)	1-2 gal/100 gal water	Up to day of harvest	4	?	
SuffOil-X (Petroleum oil)	1-2 gal/100 gal water	Up to day of harvest	4	?	Do not mix with sulfur products.
OTHER					
SucraShield (Sucrose octanoate esters)	0.8-1% vol to vol solution	0	48	?	Use between 25 and 400 gal per acre of mix per acre depending on type, growth state and spacing of crop.

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.
PHI = pre-harvest interval, REI = restricted-entry interval.

13.3 Seedcorn Maggot, *Delia platura*

Time for concern: Mid-May through late June

Key characteristics: Adult flies resemble small house flies, are slender, 1/4 inch long, and grayish black in color. Maggots are yellowish white and infest seeds and other below-ground plant parts. See Cornell [fact sheet](#) for photo (Link 52). Maggots are only a problem from planting to plant emergence. Damaged plants are weak, may not develop, or may be delayed in maturity. Stand may be poor. See Cornell [photo of damage](#) (Link 53) or see a Cornell bulletin with [more information about damage](#) (Link 54).

Management Option	Recommendations for Seedcorn Maggot
Scouting/thresholds	Although there are multiple generations per year, the first generation is the important one. No thresholds are available. Five maggots per seed are required to significantly reduce stands of snap beans.
Resistant varieties	Snap beans are particularly susceptible.
Planting date	Planting after the first generation maggots have pupated will reduce damage. In New York, this occurs around June 21. Additionally, plantings after this period are less susceptible to maggot damage because warmer soil temperatures allow seedlings to emerge soon after planting, reducing the adult egg-laying period and young larval feeding period.
Site selection/preparation	Seedcorn maggots prefer soil with high organic matter. Incorporate crop residues at least 2 weeks before planting. Using shallow planting and other means to speed up germination and emergence will reduce damage.

Seed selection/treatment	No seed treatments are currently approved for organic production.
Natural enemies	Natural enemies can be preserved by using pesticides that are less harmful to them. Use Reference 5 see Cornell guide to natural enemies (Link 47).

13.4 European Corn Borer (ECB), *Ostrinia nubilalis*

Time for concern: June through September.

Key characteristics: The eggs are laid in fish scale-like masses on the undersides of leaves. Larvae are cream colored and have a relatively smooth appearance with rows of brown spots. The head capsule is dark brown, and mature larvae are 3/4 inch long. Pupae are reddish brown, but the bean crop is harvested before pupation occurs. Adults are light yellow with reddish brown markings and are about one inch in length. See Cornell [fact sheet](#) for photo (Link 55) or Reference 3. European corn borer feeding will damage pods and their feeding within the pod will cause contamination at harvest. See Cornell [photo of damage](#) (Link 56). See reference 15.

Management Option	Recommendations for European Corn Borer
Scouting/thresholds	There are no formal thresholds for this pest on snap bean. Because tolerance for this pest is extremely low, insecticide use is relied upon heavily. Decision for control should be based on the stage of the bean crop and level of moth activity. Beans should be treated only when they are in the vulnerable stage, i.e., from early bud until early pod development. Moth activity can be monitored using traps (black light or pheromone) and trap catch information is updated on the Timely Tips section of the following website: Penn State PestWatch website (Link 57).
Natural enemies	A variety of natural enemies help suppress ECB populations including predatory lady beetles, minute pirate bugs and lacewings, and fly and wasp parasitoids. Natural enemies can be preserved using pesticides that are less harmful to them. Use Reference 5 or see the Cornell publication Biological Control: A Guide to Natural Enemies in North America (Link 47). Trichogramma ostriniae, a parasitoid of eggs, has been used successfully for ECB control in sweet corn and is commercially available. Releases should be timed to susceptible crop stage and ECB flight peaks.
Resistant varieties	No resistant varieties are available.
Crop rotation	This is not a viable management option
Site selection	Avoid planting in or adjacent to a field that was planted with corn the previous year. However, this is likely applicable only to early-planted fields. Late-planted snap bean fields adjacent to mature untreated corn could be at risk for attack by second-generation corn borers that emerge from the mature corn.
Postharvest and Sanitation	These are not currently viable management options

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide’s effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS](#)) website (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

13.4 Labeled Products for Management of European Corn Borer					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
BIOLOGICALS (<i>Bacillus thuringiensis</i> subsp. <i>Kurstaki</i>)					
Deliver	0.25-1.5 lbs	0	4	3	<i>Bacillus thuringiensis</i> products not effective in 2/2 trials. Must be eaten by the larvae to be effective; most effective against small, newly hatched larvae, so good scouting program to determine early infestations is recommended.
Javelin WG	0.12-1.5 lbs	0	4	3	
BIOLOGICALS (Spinosad)					
Entrust	1.25-2 oz	3	4	1	Effective in 41/47 trials. 33 studies showed good control of caterpillars, including ECB, Organic Resource Guide
BOTANICAL					
Aza-Direct (Azadirachtin)	1-2 pints	0	4	1	Can use for ECB, but requires an intensive scouting program.
Pyganic Crop Protection 1.4 EC (Pyrethrin)	16-64 oz	0	12	?	

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.
PHI = pre-harvest interval, REI = restricted-entry interval.

13.5 Two-Spotted Spider Mite, *Tetranychus urticae*

Time for concern: June through September.

Key characteristics: The adult mite is yellow to dark green with two or four black, dorsal spots. See Cornell [fact sheet](#) (Link 58). Heavy damage may cause leaves to drop. Hot, dry weather favors spider mite outbreaks.

Management Option	Recommendation for Two-Spotted Spider Mite
Natural enemies	Natural enemies help to control spider mite populations. See the Cornell guide Biological Control: A Guide to Natural Enemies in North America (Link 47) or use Reference 5.
Resistant varieties	No resistant varieties are available.
Crop rotation, Site selection, and Sanitation, Postharvest	These are not currently viable management options.
Note(s)	Dry, warm conditions may increase the chance of a two-spotted spider mite infestation.

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide’s effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)) (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 13.5 Pesticides Labeled for Management of Two-Spotted Spider Mites					
Class of Compounds Product Name (active ingredient)	Product Rate/Acre	PHI (hours)	REI (hours)	Efficacy	Comments
BOTANICALS					
Azahar (Azadirachtin)	12-41 fl oz	0	4	?	
AzaMax (Azadirachtin)	1.33 fl oz/1000 ft ²	0	4	?	
Safer Brand #567 Pyrethrin & Insecticidal Soap Concentrate II (Pyrethrin & potassium salts of fatty acids)	1 gal spray mix/700 ft ² plant surface area	-	12	?	See label for specific mixing instructions.
Trilogy (Neem extract)	0.5-2.0% in 25-100 gal water/A	None listed	4	?	Maximum labeled use of 2 gal/acre/application.
OILS					
Glacial Spray Fluid (Mineral oil)	0.75-1 gal/100gal	Up to day of harvest	4	?	See label for specific application volumes
Golden Pest Spray Oil (Soybean oil)	2 gal	-	4	?	
Organic JMS Stylet Oil (Paraffinic oil)	3-6 qt/100 gal	0	4	?	See label for restrictions when combined with sulfur.
Saf-T-Side (Petroleum oil)	1-2 gal/100 gal water	Up to day of harvest	4	?	
SuffOil-X (Petroleum oil)	1-2 gal/100 gal water	Up to day of harvest	4	?	Do not mix with sulfur products.
OTHER					
SucraShield (Sucrose octanoate esters)	0.8-1% vol to vol solution	0	48	?	Use between 25 and 400 gal per acre of mix per acre depending on type, growth state and spacing of crop.

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.
PHI = pre-harvest interval, REI = restricted-entry interval.

13.6 Tarnished Plant Bug (TPB), *Lygus lineolaris*

Time for concern: June through September

Key characteristics: The adult is an oval, brown bug, mottled with various shades of reddish and yellowish brown, and about 1/4 inch long. The eggs are elongated and curved with a square outer end. Nymphs are small and greenish yellow. See Cornell [fact sheet](#) (Link 59). The nymphs feed more than the adults. The sucking injury from the nymphs causes buds to drop, pods to be misshapen, and plants to be stunted and distorted. Tarnished plant bug can be a problem from bloom through harvest. See Cornell [photo of damage](#) (Link 60) or Reference 14.

Management Option	Recommendations for Tarnished Plant Bug
Scouting/thresholds	Check for TPB on pigweed seed heads or by using a sweep net. Trials on snap beans indicated no yield reductions occurred with five adult TPBs or less per plant at blossom through pin pod stages.
Natural enemies	Natural enemies that can help control TPB populations can be preserved by using pesticides that are less harmful to them. See the Cornell guide <i>Biological Control: A Guide to Natural Enemies in North America</i> (Link 47) or use Reference 5.

Resistant varieties	No resistant varieties are available. Although the TPB will feed on bean pods under New York's growing conditions, most snap bean varieties are somewhat tolerant with minimal pod-spotting resulting at harvest. It would be advisable to schedule plantings of susceptible varieties for early-season harvests since TPB numbers increase in bean fields as the season progresses.
Crop rotation, Site selection, Postharvest, and Sanitation	Effective weed management could eliminate potential hosts for tarnished plant bugs in fields and thereby minimize risk of injury to the snap bean crop. TPB has a wide-range of hosts including many weeds and is particularly attracted to flower buds. Minimizing weeds that tend to bloom prior to bean blooms can help reduce the overall population of TPB in the field. Avoid situations in which snap beans, in a vulnerable stage, are near hayfields where TPB numbers may build up and move into beans when the hay is cut.

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 13.6 Pesticides Labeled for Management of Tarnished Plant Bug					
Class of Compounds Product Name (active ingredient)	Product Rate/Acre	PHI (hours)	REI (hours)	Efficacy	Comments
BOTANICAL					
Azahar (Azadirachtin)	12-41 fl oz	0	4	?	
AzaMax (Azadirachtin)	1.33 fl oz/1000 ft ²	0	4	?	
Safer Brand #567 Pyrethrin & Insecticidal Soap Concentrate II (Pyrethrin & potassium salts of fatty acids)	1 gal spray mix/700 ft ² plant surface area	-	12	?	See label for specific mixing instructions.
OILS					
Golden Pest Spray Oil (Soybean oil)	2 gal	-	4	?	Only for use against larvae.

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.
PHI = pre-harvest interval, REI = restricted-entry interval.

14. SLUGS

Time of concern: May through September.

Key characteristics: Adult slugs are between one and two inches in length. See Ohio State University [fact sheet](#) (Link 61). Slugs can overwinter at any stage of development. Although slugs cannot survive prolonged subzero temperatures or desiccation, the burrows of small mammals and worms provide insulation from the weather. Slugs begin to move, hatch, feed, and lay eggs in the spring when temperatures are consistently above 40°F. There is often little or no slug activity in the field during periods of dry weather; however, there may be extensive feeding when the weather is damp.

Management Option	Recommendations for Management of Slugs
Scouting/thresholds	Record the occurrence and severity of slug damage. No thresholds have been established.
Resistant varieties	No resistant varieties are available.
Cultural	Practices that help dry the soil surface for example conventional tillage, good weed control, and using raised beds that dry out more readily than flat beds, will reduce slug populations. Heavy organic mulch creates an ideal environment for slugs.

At the time this guide was produced, the following materials were labeled in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide’s effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. Those pesticides meeting requirements in EPA Ruling 40 CFR Part 152.25(b) (also known as 25(b) pesticides) do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 14. Pesticides Labeled for Management of Slugs					
Class of Compound Product (Active ingredient)	Rate/A	PHI (Days)	REI (hours)	Efficacy	Comments
IRON					
<i>Sluggo AG</i> Iron phosphate	20-44 lbs.	0	0	?	Treat field perimeter. Lasts up to 4 weeks.

Efficacy: 1-effective in some research studies, 2- mixed efficacy results, 3-not effective, ?- not reviewed or research not available.
PHI = pre-harvest interval, REI = restricted-entry interval.

15. PESTICIDES & ABBREVIATIONS MENTIONED IN THIS PUBLICATION

Table 1. Insecticides and molluscicide mentioned in this publication

TRADE NAME	COMMON NAME	EPA REG. NO.
Aza-Direct	<i>Azadirachtin</i>	71908-1-10163
AzaGuard	<i>Azadirachtin</i>	70299-17
Azahar	<i>Azadirachtin</i>	71908-1-10163
AzaMax	<i>Azadirachtin</i>	71908-1-81268
Deliver	<i>Bt var. kurstaki</i>	70051-69
Entrust	<i>spinosad</i>	62719-282
Glacial Spray Fluid	<i>Mineral oil</i>	34704-849
Golden Pest Spray Oil	<i>Soybean oil</i>	57538-11
Javelin WG	<i>Bt var. kurstaki</i>	70051-66
Neemazad 1%EC	<i>Azadirachtin</i>	70051-104
Neemix 4.5	<i>Azadirachtin (neem)</i>	70051-9
Organic JMS Stylet Oil	<i>Paraffinic oil</i>	65564-1
Pyganic Crop Protection EC 1.4	<i>pyrethrin</i>	1021-1771
Pyganic Crop Protection EC 5.0	<i>pyrethrin</i>	1021-1772
Safer Brand #567	<i>Pyrethrin & Potassium salts of fatty acids</i>	59913-9
Saf-T-Side	<i>Petroleum oil</i>	48813-1
SucraShield	<i>Sucrose octanoate esters</i>	70950-2-84710
SuffOil-X	<i>Petroleum oil</i>	48813-1-68539
Sluggo Ag	<i>iron phosphate</i>	67702-3-54705
Trilogy	<i>Neem extract</i>	70051-2

Table 2. Fungicides mentioned in this publication

TRADE NAME	COMMON NAME	EPA REG. NO.
ActinoGrow	<i>Streptomyces lydicus</i>	73314-1
Actino-Iron	<i>Streptomyces lydicus</i>	73314-2
Actinovate AG	<i>Streptomyces lydicus</i>	73314-1
Basic Copper 53	<i>basic copper sulfite</i>	45002-8
Champ WG	<i>Copper hydroxide</i>	55146-1
Cueva Fungicide Concentrate	<i>Copper hydroxide</i>	67702-2
Contans WG	<i>Coniothyrium minitans</i>	72444-1
Heads Up Plant Protection	<i>Chenopodium quinoa</i>	81853-1
MicroSulf	<i>sulfur</i>	55146-75
Microthiol Dispers	<i>Sulfur</i>	70506-187
Milstop	<i>Potassium bicarbonate</i>	70870-1-68539
Mycostop	<i>Streptomyces griseoviridis</i>	64137-5
Mycostop Mix	<i>Streptomyces griseoviridis</i>	64137-9
Nu Cop 50 WP	<i>cupric hydroxide</i>	45002-7
Organic JMS Stylet Oil	<i>Paraffinic oil</i>	65564-1
Serenade ASO	<i>Bacillus subtilis</i>	69592-12
Serenade MAX	<i>Bacillus subtilis</i>	69592-11
Sporan	<i>rosemary oil, clove oil, thyme oil</i>	exempt – 25(b)
T-22 HC	<i>Trichoderma harzianum</i>	68539-4
Trilogy	<i>neem oil</i>	70051-2

Abbreviations and Symbols Used in This Publication

A	acre	NE	not effective
AG	agricultural use label	NI	no information
AR	annual rye	NFT	not frost tolerant
ASO	aqueous suspension-organic	P	phosphorus
AS	aqueous suspension	PHI	pre-harvest interval
DF	dry flowable	P ₂ O ₅	phosphorus oxide
EC	emulsifiable concentrate	PR	perennial rye
F	flowable	R	resistant varieties
HC	high concentrate	REI	reentry interval
K	potassium	WP	wettable powder
K ₂ O	potassium oxide	WG	water dispersible granular
N	nitrogen	WPS	worker protection standards

16. REFERENCES

- Babadoost, M. University of Illinois. (2006). *New Strategies for Management of Vegetable Diseases in Organic and Traditional Farms*. (http://www.sare.org/reporting/report_viewer.asp?pn=LNC03-228).
- Caldwell, B. Rosen, E. B., Sideman, E., Shelton, A. M., Smart, C. (2005). *Resource Guide for Organic Insect and Disease Management*. New York State Agricultural Experiment Station, Geneva, NY. (<http://www.nysaes.cornell.edu/pp/resourceguide/>).
- Colorado State University. (1996). *Dry Bean Production and Pest Management*. Regional Bulletin 562A, Cooperative Extension Resource Center, 115 General Services Building, Fort Collins, CO.
- Hall, R. ed. (1991). *Compendium of Bean Diseases*. (73 pp). American Phytopathologic Society Press.
- Hoffmann, M. P., and Frodsham A. C. (1993). *Natural Enemies of Vegetable Insect Pests*. (64 pp). Cornell Cooperative Extension. New York State Agricultural Experiment Station, Geneva, NY.
- Iowa State University. (1983). *Recognition and Management of Dry Bean Production Problems*. North Central Regional Extension Publication 198. Cooperative Extension Resource Center, Ames, IA.
- New York State Integrated Pest Management Program. (2008). *Integrated Crop and Pest Management Guidelines for Commercial Vegetable Production*: Chapter 13, Beans-Dry and Snap. Cornell Cooperative Extension, Geneva, NY. (<http://www.nysaes.cornell.edu/recommends/13frameset.html>).
- Pennsylvania State University. (1987). *Weed identification* (pp. 1-32). College of Agriculture, Cooperative Extension Service, University Park, PA.
- Petzoldt, C. H., Pederson L. H., and Koplinka-Loehr, C. eds. (1990). *Snap Bean Pest Management: A Guide to Regular Field Monitoring in New York*. IPM Publication. 105b. New York State Agricultural Experiment Station, Geneva, NY.
- Rosen, C. J., Bierman, P. M. (2005). *Using Manure and Compost as Nutrient Sources for Fruit and Vegetable Crops*. University of Minnesota, MN, (<http://extension.umn.edu/distribution/horticulture/M1192.html>).
- Sarrantonio, M. (1994) *Northeast Cover Crop Handbook*. Rodale Institute, PA. (<http://www.rodaleinstitutesstore.org/store/customer/product.php?productid=783&cat=460&page=1>).
- Wilsey, W.T., Weeden, C.R., Shelton, A.M., *Pests in the Northeastern United States*. (<http://www.nysaes.cornell.edu/ent/factsheets/>).
- Zitter, T. A., McGrath, M. T. *Vegetable MD Online*. (http://vegetablemdonline.ppath.cornell.edu/factsheets/Bean_List.htm).
- Muka, A. A. (1983). Tarnished plant bug, In *Vegetable Crops: Insects of Bean, Potato, and Celery* (p. 771.00). New York State Agricultural Experiment Station, Geneva, NY.
- Showers, W. B., Witkowski J. F., Mason C. E., Calvin D. D., Higgins R. A., and Dively G. P. (1989). *European Corn Borer: Development and Management*. North Central Regional Extension Publication 327. Iowa State University, Ames, IA.
- Tingey, W. M., Muka A. A. (1983). Potato leafhopper, In *Vegetable Crops: Insects of Vegetables* (p. 760.20). New York State Agricultural Experiment Station, Geneva, NY.
- Stivers, L.J., Brainard, D.C. Abawi, G.S., Wolfe, D.W. (1999) *Cover Crops for Vegetable Production in the Northeast*. Cornell Cooperative Extension, Ithaca, NY (<http://ecommons.library.cornell.edu/bitstream/1813/3303/2/Cover%20Crops.pdf>).

- 18 Dillard, H. R., and Cobb, A. C. 2008. *Alternaria alternata* and *Plectosporium tabacinum* on snap beans: Pathogenicity, cultivar reaction, and fungicide efficacy. Online. Plant Health Progress doi:10.1094/PHP-2008-1212-01-RS.

17. WORLD WIDE WEB LINKS

All links accessed 13 January 2009

General

- 1 Hardiness Zone Map for New York (<http://www.gardening.cornell.edu/weather/images/zonclg.jpg>).
- 2 Pesticide Product Ingredient, and Manufacturer System (PIMS). (<http://pims.psur.cornell.edu>).
- 3 The Network for Environment and Weather Awareness (NEWA). New York State Integrated Pest Management Program. (<http://newa.nysaes.cornell.edu/public/default.htm>).
- 4 Cornell University. New York State Integrated Pest Management Program. (2002). *Elements of IPM for Snap Beans in New York State*. (<http://nysipm.cornell.edu/elements/snapbean.asp>).

Certification

- 5 Organic Materials Review Institute. (<http://www.omri.org/>).
- 6 New York Department of Agriculture and Markets, Organizations Providing Organic Certification Services for Producers and Processors in New York State. (<http://www.agmkt.state.ny.us/AP/organic/docs/Organizations%20Providing%20Organic%20Certification%20Services%20for%20Farmers%20and%20Processors%20in%20New%20York%20State.pdf>).
- 7 New York Department of Agriculture and Markets, Organic Farming Resource Center. (<http://www.agmkt.state.ny.us/AP/organic/>).
- 8 Agriculture Marketing Service, National Organic Program. (<http://www.ams.usda.gov/nop/NOP/standards/ProdHandPre.html>).
- 9 National Organic Program Final Rule 2000. (<http://www.ams.usda.gov/AMSV1.0/ams.fetchTemplateData.do?template=TemplateF&navID=RegulationsNOPNationalOrganicProgramHome&rightNav1=RegulationsNOPNationalOrganicProgramHome&topNav=&leftNav=NationalOrganicProgram&page=NOPRegulations&resultType=&ac>).
- 10 National Sustainable Agriculture Information Service, Organic Farming. (<http://attra.ncat.org/organic.html>).
- 11 Rodale Institute. (<http://www.rodaleinstitute.org/>).

Soil Health, Cover Crops, and Crop Rotation

- 12 Björkman, Thomas. Cornell University, *Cover Crops for Vegetable Growers*. (<http://www.nysaes.cornell.edu/hort/faculty/bjorkman/covercrops/decisiontool.php>).
- 13 Magdoff, F., Van Es, H., (2000). Sustainable Agriculture Research and Education, *Building Soils for Better Crops, 2nd Edition*. (<http://www.sare.org/publications/bsbc/index.htm>),
- 14 Cornell University, Department of Horticulture. *Soil Health Website*. (<http://www.hort.cornell.edu/soilhealth/extension/test.htm>).
- 14 a Mohler, C. L. and Johnson, S. E., editors. (2009). *Crop Rotation on Organic Farms: A Planning Manual*. Sustainable Agriculture Research and Education. Natural Resource, Agriculture and Engineering Service. Cooperative Extension, Ithaca NY. (http://www.nraes.org/nra_crof.html).

Weed Management

- 15 Bowman, G., (1997). The Sustainable Agriculture Network. *Steel in the Field*. Beltsville, MD. (<http://www.sare.org/publications/steel/index.htm>).
- 16 Cornell University, Weed Ecology and Management Laboratory. (<http://www.css.cornell.edu/weedeco/>).
- 17 Rutgers University, *New Jersey Weed Gallery* (<http://njaes.rutgers.edu/weeds/>).
- 18 University of Vermont, Videos for Vegetable and Berry Growers. (<http://www.uvm.edu/vtvegandberry/Videos/videos.html>).
- 19 Sullivan, P., National Sustainable Agriculture Information Service (formerly ATTRA), *Principles of Sustainable Weed Management for Croplands*. (<http://attra.ncat.org/attra-pub/weed.html>).
- 20 Colquhoun, J., Bellinder, R., Cornell University. *New Cultivation Tools for Mechanical Weed Control in Vegetables*. (<http://www.vegetables.cornell.edu/pubs/newcultivationmech.pdf>).

Crop and Soil Nutrition

- 21 Cornell Nutrient Analysis Laboratory. (<http://cnal.cals.cornell.edu/>).
- 22 Agri Analysis, Inc.. (<http://www.agrianalysis.com/>).
- 23 A&L Eastern Agricultural Laboratories, Inc. (<http://al-labs-eastern.com/>).
- 24 The Pennsylvania State University, Agricultural Analytical Services Laboratory. (<http://aasl.psu.edu>).
- 25 Cornell University, The Dairy One Forage Lab, Ithaca, NY. (<http://www.dairyone.com/forage/default.asp>).
- 26 University of Massachusetts, Soil and Plant Tissue Testing Laboratory. (<http://www.umass.edu/plsoils/soiltest/>).
- 27 Analytical Laboratory and Maine Soil Testing Service, University of Maine. (<http://anlab.umesci.maine.edu/>).
- 28 Rosen, C., Bierman, P. Using Manure and Compost as Nutrient Sources for Fruit and Vegetable Crops. University of Minnesota. (<http://www.extension.umn.edu/distribution/horticulture/M1192.html>).
- 28 a. The Pennsylvania State University. (2007-08). *Penn State Agronomy Guide*. Department of Agronomy. University Park, PA

Managing Bean Diseases

- 29 Cornell University, Vegetable MD Online. (2008) *Green Bean: Disease Resistance Table*. (http://vegetablemdonline.ppath.cornell.edu/Tables/Bean_GreenTable.html).
- 30 Cornell University, Vegetable MD Online. (2008) *Wax Bean: Disease Resistance Table*. (http://vegetablemdonline.ppath.cornell.edu/Tables/Bean_WaxTable.html).
- 31 Abawi, G. S., and Hunter J. E. (1979). **White Mold of Beans in New York**. (4 pp.). New York's Food and Life Sciences Bulletin 77. New York State Agricultural Experiment Station, Geneva, NY. (<http://ecommons.library.cornell.edu/bitstream/1813/5083/1/FLS-077.pdf>).
- 32 Vegetable MD online. *Photo Gallery of Important New York Vegetable diseases: Bean: White Mold*. (http://vegetablemdonline.ppath.cornell.edu/photopages/impt_diseases/beans/Bean_White.htm).
- 33 George S. Abawi and J.E. Hunter (1979). Vegetable MD Online. **White Mold of Beans**. New York's Food and Life Sciences Bulletin No.77 (http://vegetablemdonline.ppath.cornell.edu/factsheets/Beans_WhiteMold.htm).
- 34 Vegetable MD online. Photo Gallery of Important New York Vegetable diseases: *Bean: Gray mold*. (http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Impt_Diseases/Beans/Bean_Gray.htm).
- 35 Abawi, G. S., Crosier D. C., and Cobb A. C. (1985). **Root Rot of Snap Beans in New York**. (8 pp). New York's Food and Life Sciences Bulletin 110. New York State Agricultural Experiment Station, Geneva, NY. (<http://www.nysaes.cornell.edu/pubs/fls/OCRPDF/110.pdf>).
- 36 Cornell University, Vegetable MD Online. *Photo Gallery of Important New York Vegetable Diseases: Bean: Pythium Root Rot* (http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Impt_Diseases/Beans/Bean_Pythium.htm).
- 37 Cornell University, Vegetable MD Online. *Photo Gallery of Important New York Vegetable diseases: Bean: Rhizoctonia Root Rot*. (http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Impt_Diseases/Beans/Bean_Rhizo.htm).
- 38 Dillard, H. R., and Legard, D. E. (1991). Cornell University, Vegetable MD Online. **Bacterial Diseases of Beans**. (p. 729.50). New York State Agricultural Experiment Station, Geneva, NY. (http://vegetablemdonline.ppath.cornell.edu/factsheets/Beans_Bacterial.htm).
- 39 Zitter, T.A., Provvidenti, R. (1984). Cornell University, Vegetable MD Online. **Virus Diseases of Snap and Dry Beans**. (http://vegetablemdonline.ppath.cornell.edu/factsheets/Virus_Beans.htm).
- 40 Cornell University, Vegetable MD Online. *Bean Virus Photo Collection*. (<http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Bean/Viruses/BVphotoList.htm>).
- 41 Zitter, T. A. (2001). Cornell UniversityVegetable MD Online. *A Checklist of Major Weeds and Crops as Natural Hosts for Plant Viruses in the Northeast*. (<http://vegetablemdonline.ppath.cornell.edu/Tables/WeedHostTable.html>).
- 42 Vegetable MD online. Photo Gallery of Important New York Vegetable Diseases: **Bean Rust**. (http://vegetablemdonline.ppath.cornell.edu/photopages/impt_diseases/beans/bean_rust.htm).
- 43 Mercure, P. S. (1998). University of Connecticut. **Bean Rust**. (<http://www.hort.uconn.edu/lpm/veg/htms/bnrust.htm>).

Managing Nematodes

- 44 Plant Disease Diagnostic Clinic. Cornell University. (<http://plantclinic.cornell.edu/fees.htm> - Nematode%20analysis).

- 45 Abawi, F.S., Gugino, B.K. (2007) Cornell University, New York State Agricultural Experiment Station. *Soil Sampling for Plant-Parasitic Nematode Assessment*. (<http://www.nysaes.cornell.edu/recommends/Nemasoilsample.pdf>).
- 46 Gugino, B.K., Ludwig, J.W., Abawi, G.S., Cornell University, New York State Agricultural Experiment Station. *A Soil Bioassay for the Visual Assessment of Soil Infestations of Lesion Nematode*. (<http://www.nysaes.cornell.edu/recommends/Lesionnemahowto.pdf>).

Managing Bean Insects

- 47 Weeden, C.R., Shelton, A.M., Hoffmann, M. P., *Biological Control: A Guide to Natural Enemies in North America*. (<http://www.nysaes.cornell.edu/ent/biocontrol/index.html>).
- 48 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Mexican Bean Beetle**. (<http://www.nysaes.cornell.edu/ent/factsheets/pests/mbb.html>).
- 49 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2001). Cornell University. Pests in the Northeastern United States. **Mexican Bean Beetle- Damage to Bean**. (http://www.nysaes.cornell.edu/ent/factsheets/damage/mbb_beans.html).
- 50 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2001). Cornell University. Pests in the Northeastern United States. **Potato Leafhopper Life Cycle**. (<http://www.nysaes.cornell.edu/ent/factsheets/pests/plh.html>).
- 51 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2001). Cornell University. Pests in the Northeastern United States. **Potato Leafhopper Damage to Beans**. (http://www.nysaes.cornell.edu/ent/factsheets/damage/plh_beans.html).
- 52 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Seedcorn Maggot Life Cycle**. (<http://www.nysaes.cornell.edu/ent/factsheets/pests/scm.html>).
- 53 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Seedcorn Maggot Damage to Beans**. (http://www.nysaes.cornell.edu/ent/factsheets/damage/scm_beans.html).
- 54 Veal, E. V., Webb D. R., and Eckenrode C. J. (1975). **Seedcorn Maggot Injury** (4 pp.). New York's Food and Life Sciences Bulletin 55. New York State Agricultural Experiment Station, Geneva, NY. (<http://www.nysaes.cornell.edu/pubs/fls/OCRPDF/55.pdf>).
- 55 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **European Corn Borer Life Cycle**. (<http://www.nysaes.cornell.edu/ent/factsheets/pests/ecb.html>).
- 56 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **European Corn Borer Damage to Bean**. (http://www.nysaes.cornell.edu/ent/factsheets/damage/ecb_beans.html).
- 57 Fleischer, S. J., Penn State, **PestWatch**. University Park, PA (<http://www.pestwatch.psu.edu/>).
- 58 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Twospotted Spider Mite Life Cycle**. (<http://www.nysaes.cornell.edu/ent/factsheets/pests/tsm.html>).
- 59 Cornell Cooperative Extension. New York State Integrated Pest Management Program. (1991). **Tarnished Plant Bug**. Insect Identification Sheet No. 21, 1991 (<http://www.nysipm.cornell.edu/factsheets/treefruit/pests/tpb/tpb.asp>).
- 60 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Tarnished Plant Bug Damage to Beans**. (http://www.nysaes.cornell.edu/ent/factsheets/damage/tpb_beans.html).
- 61 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Slugs Life Cycle**. (<http://ohioline.osu.edu/hyg-fact/2000/2010.html>).

Pesticide Use, Safety, and Sprayer Calibration

- 62 Cornell Integrated Crop and Pest Management Guidelines (2009). *Chapter 6 Pesticide Information and Safety*. (<http://www.nysaes.cornell.edu/recommends/6frameset.html>).
- 63 *Calibration: Backpack Sprayer*. Pesticide Environmental Stewardship. Center for Integrated Pest Management. (<http://pesticidestewardship.org/calibration/Pages/BackpackSprayer.aspx>)
- 64 Dill, J. and Koehler, G. (2005). Agricultural Pocket Pesticide Calibration Guide. University of Maine Cooperative Extension and Northeastern IPM Center. (<http://pronewengland.org/INFO/PROpubs/CalibrationGuide-small.pdf>).
- 65 Landers, A., Knapsack Sprayers: General Guidelines for Use. Cornell University, Ithaca, N.Y. (http://www.nysaes.cornell.edu/ent/faculty/landers/pestapp/publications/veg/knapsack_sprayer.doc).
- 66 Miller, A. and Bellinder, R. (2001) Herbicide Application Using a Knapsack Sprayer. Department of Horticultural Science, Cornell University, Ithaca, N.Y. (<http://www.hort.cornell.edu/spray/southasia/pdfs/knapsack.pdf>).
- 67 Federal Insecticide Fungicide Rodenticide Act (FIFRA). (2009). *Electronic Code of Federal Regulations*. Title 40: Protection of the Environment. Part 152 Pesticide registration and classification procedures. Subpart B- exemptions. Part 152.25(b). (<http://ecfr.gpoaccess.gov/cgi/t/text/text-idx?c=ecfr&sid=26e5b67c614d04a9a5b7db5acea29a62&rgn=div8&view=text&node=40:23.0.1.1.3.2.1.2&idno=40>).

ORGANIC SNAP BEAN PRODUCTION

- 68 Office of prevention, pesticides and toxic substances. (2009). *Inert ingredients eligible for FIFRA 25(b) pesticide products*. United States Environmental Protection Agency. Washington DC. (http://www.epa.gov/opprd001/inerts/section25b_inerts.pdf).
- 69 Federal Insecticide Fungicide Rodenticide Act (FIFRA). (2009). *Electronic Code of Federal Regulations*. Title 7: Agriculture. National Organic Program, Part 205, sections 600-606. (<http://ecfr.gpoaccess.gov/cgi/t/text/text-idx?type=simple;c=ecfr;cc=ecfr;sid=4163ddc3518c1ffdc539675aed8efe33;region=DIV1;q1=national%20organic%20program;rgn=div5;view=text;idno=7;node=7%3A3.1.1.9.31>).

This guide is published by the New York State Integrated Pest Management Program, which is funded through Cornell University, Cornell Cooperative Extension, the New York State Department of Agriculture and Markets, the New York State Department of Environmental Conservation, and USDA-NIFA. Cornell Cooperative Extension provides equal program and employment opportunities. NYS IPM Publication number 132, version 2. February 2011 . www.nysipm.cornell.edu.